

## Effect of Drought on Qualitative and Quantitative Parameters of Zahedan Plain Aquifer

Masome Sarhadi<sup>1</sup>, Mohammad Nohtani<sup>2\*</sup> and Mohammad Reiki<sup>3</sup>

*1 Former Master Student, Department of Rangeland and Watershed Management, University of Zabol, Zabol, Iran*

*2 Assistant Professor, Department of Rangeland and Watershed Management, Faculty of Water and Soil, University of Zabol, Zabol, Iran*

*3 Former Master Student, Department of Rangeland and Watershed Management, University of Zabol, Capacity Building for the International Project of MENARID in Sistan and Baluchistan Province, Iran*

Received: 2 May 2016 / Accepted: 15 August 2016 / Published Online: 9 December 2016

**ABSTRACT** The effect of drought indices on hydrostatic variations and qualitative parameters of the Zahedan Plain's aquifer was investigated. For this purpose, 12-year statistics from observational and piezometry wells in the plain, the drought index of percentage normal (PN), standardized precipitation index (SPI), and 30-year precipitation statistics of Zahedan station were used. Results revealed that the PN index had a greater variety than SPI. The aquifer hydrograph showed a rise of 1.37 m in water level from the year 2002-2003 to 2013-2014. Considering the drought conditions in the plain, this rise in the water level can be attributed to the reduced water extraction from the aquifer, because the water requirement of Zahedan city was supplied from other sources, viz. Chah-nimehs. Also, there was a significant correlation between the parameters of anion, TDS, and Na<sup>+</sup> with SPI. However, PN index did not have any significant relationship with the quality parameters. There was a significant correlation between water level balance and SPI only at the level of 1%.

**Key words:** *Acquirer's unit hydrograph, Drought indices, Ground water quality, Zahedan plain*

### 1 INTRODUCTION

Evaluation of ground water resources and their stability in arid and semi- arid regions are of great importance, as they are usually the sole water supplying source for drinking and agricultural purposes (Nosrati and Eeckhaut, 2011). Precipitation permeates into soil and upon reaching impermeable layers, it develops ground water reserves. In recent years, drought has resulted in a considerable decrease in surface waters, while use of ground waters has increased, which has led to both dramatic decrease in the level of ground water acquirers

and changes in their quality (Afzali and Shahedi, 2014).

Despite the fact that ground water resources are exploited in different areas, a strong relationship between the indicator SPI and ground water levels exist in many cases, which was used to relate the efficiency of ground water levels in Australia with its major drought pattern (Khan *et al.* 2008). Ahmadvash *et al.* (2013) examined the effects of recent droughts on ground water resources in Azarshahr Plain. They found increase in ground water level by 0.59 m from June 2002 to June 2010, and

\* Corresponding author: Assistant Professor, Faculty of Water and Soil, University of Zabol, Zabol, Iran, Tel: +98 911 9648185, Email: m.nohtani@uoz.ac.ir

Also increased electrical conductivity, chloride and sodium concentrations due to the intrusion of the salty water from Urmia Lake. Watmough and Juckers (2014) examined the effects of simulated drought and found the level of aquifer decreased, but the effective water chemical factors increased. In general, the majority of studies reveal a descending ground water hydrostatic level and effective qualitative factors in the long term drought. Najafzadeh *et al.* (2015) found that the water level and quality factors of the ground water had dropped in much of the Mah-Velayat Plain as the result of drought. Application of standardized precipitation index (SPI) in Kashan (Aleboali *et al.*, 2016) and Shahre-Babak plains (Jahanshahi *et al.*, 2016) showed a significant relationship between the ground water level and drought.

The experience of drought in recent years in Sistan-Baluchistan and the importance of the ground water resources in this province highlight the significance of examination of drought in this region. In this research, the SPI


index was used to assess the impact of drought on the ground water level and qualitative parameters in Zahedan Plain aquifer.

The findings of this research can help planners and managers in making decisions in line with variations in climatic elements.

## 2 MATERIALS AND METHODS

### 2.1 Study region

Zahedan Plain is situated in Sistan-Baluchistan Province ( $60^{\circ} 39' - 60^{\circ} 56' E$  and  $29^{\circ} 23' - 29^{\circ} 35' N$ ). Considering the distribution of wells, one major aquifer with an area around  $210 \text{ km}^2$  in alluvial sediments was considered for this study (Figure 1). The average height of mountain and plain this region is 1686 and 1454 meters, respectively. The direction of the ground water in the aquifer is from west to east and in the southern region is northwards, eventually leaving the aquifer from the northeast (Regional Water Bureau, 2011).



**Figure 1** Location of the study area in Sistan-Baluchistan Province (top) and distribution of wells (bottom)

### 2.2 Index of Percentage of Normal Precipitation (PN)

Being always positive and limited to zero from the bottom, PN index is has no theoretical limitation from the top. Its main concept is

division of real precipitation by normal precipitation. It is calculated this index, Equation (1), has been used (Nahvinia *et al.*, 2008).

$$PN = \left(\frac{P}{P_0}\right) \times 100 \quad (1)$$

Where P is monthly precipitation or the period of interest (mm), P<sub>0</sub> the average long-term precipitation during this statistical period. Considering the different values of PN, various drought intensities are recognizable (Table 1).

**Table 1** Classification of drought intensities

Drought Classification	Value
Normal	80-120
Slight drought	70-80
Moderate drought	55-70
Severe drought	40-55
Extreme drought	Less than 40%

### 2.3 Standardized precipitation index (SPI)

In this method, the duration of drought period is determined by the beginning and termination of negative figures of SPI, with the cumulative values of SPI representing the magnitude and intensity of drought period (McKee *et al.*, 1993). To calculate this method, Equation (2) is used.

$$SPI = \frac{(P - P_0)}{SD} \quad (2)$$

Where SPI is standardized precipitation index, P monthly or annual precipitation (mm), P<sub>0</sub> the average monthly or annual precipitation (mm), SD standard deviation within the period of interest. Following the calculation of the SPI for each period, using Table 2, the years in which drought has occurred in the region were determined (McKee *et al.*, 1993).

**Table 2** Values of classified standardized precipitation index (McKee *et al.*, 1993)

Drought Classification	Value
Extremely wet	+2
Very wet	1/5 up to 1/99
Moderately wet	1 up to 1/49
Near normal	-0/99 up to 0/99
Moderately dry	-1/49 up to -1
Severely dry	-1/99 up to -1/5
Extremely dry	Less than -2

### 2.4 Plotting unit hydrograph of the aquifer

The unit hydrograph on the level of the balance of ground waters was plotted in order to observe the trend of changes in the ground water level of the region over time. After organizing the statistics, first to generalize the measured values to the region's level, Theissen function was used in the Arc Map 9.3 software. Next, the aquifer's water balance hydrograph was plotted in Excel with the aim of achieving an overview of the trend of changes in the ground water level. The aquifer unit hydrograph was obtained by Equation (3) (Akbariet *al.*, 2009).

$$h_p(y, m) = \lambda_1 * h_{w_1}(y, m) + \lambda_2 * h_{w_2}(y, m) + \dots + \lambda_n * h_{w_n}(y, m) \quad (3)$$

Where hp (y,m) is the aquifer balance in the year y and month m. λ<sub>1</sub>, λ<sub>2</sub>, ..., λ<sub>n</sub> the weight of observational wells of w<sub>1</sub>, w<sub>2</sub>, ..., w<sub>n</sub>. h<sub>w1</sub> (y,m) the balance of observational well of w<sub>1</sub> in the year y and month m. h<sub>w<sub>n</sub></sub> (y,m) the balance of observational well of w<sub>n</sub> in the year y and month m.

### 2.5 Qualitative zoning of the aquifer

Geographical location and water qualitative data for each well were fed to GS<sup>+</sup> software to determine suitable semivariogram. The results were transferred to the ArcGIS 9.3 software, whereby based on Kriging method, the zoning maps of the qualitative status of ground water in the plain of interest were prepared (Ahmed, 2002; Arsalan, 2012; Taghizadeh Mehrjerdi, 2008).

In selecting the best model of fitted semivariogram, the indices should reach their optimal value. Greater value of R<sup>2</sup> and the lower value of residual sum of squares (RSS) represent the greater accuracy of the fitted model (Jahanshahi *et al.*, 2014; Sarhadi, 2015).

## 2.6 Water quality in terms of agriculture

Wilcox diagram, based on electrical conductivity and sodium absorption rate, was used for this purpose (Wilcox, 1955), in which the sodium absorption rate is calculated by Equation (4).

$$SAR = \frac{Na}{\sqrt{\frac{Ca+Mg}{2}}} \quad (4)$$

Where Na, Ca and Mg denote the amounts of sodium, calcium, and magnesium ions, respectively, in terms of m Eq l<sup>-1</sup> in the water sample.

## 2.7 Water quality in terms of drinking

The Schoeller diagram was used for this purpose (Schoeller, 1962), in which only typical anions and cations along with mineral properties of water are considered.

## 2.8 Determining the type of water

The type of ground water was determined using Piper's diagram (Goovaerts, 1997), in which water is classified according to its cation (viz., calcic, sodic, etc.) and anion constituents (viz., bicarbonated, sulfated, etc.).

## 3 RESULTS

### 3.1 Standardized precipitation index (SPI)

The results indicated that the region tolerated very dry conditions in 2002 and 2003, except

for Mar. and in 2004 except for the months of Jan., Feb. and Mar. The conditions were relatively normal and no sign of drought was observed in 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, and 2013. Moreover, in Mar. 2005, 2007, and 2010, the conditions were mild and humid.

### 3.2 Percent of normal index (PN)

A weak precipitation occurred in Jan. 2003, Apr. 2005, Mar. 2009, and Nov. 2010. A moderate drought happened in Feb. 2005, Jan. 2005, 2006, and 2008, Dec. 2007, and Nov. 2013. Moreover, severe droughts occurred in Apr. 2003, Jan. 2002 and 2012, and Feb. 2003 and 2008. In the other month of the year, incidents of extremely severe droughts were observed.

### 3.3 Changes in the hydrostatic level of Zahedan plain's aquifer

The unit hydrographs of the plain indicated increased level of the ground water by 1.37 m. within 12 years of the investigated period (Figure 2).

### 3.4 Quantitative parameters of ground water

Using the GS<sup>+</sup> software, the conventional Kriging method was chosen for the estimation of 14 quantitative parameters in 2002-2003 and 2012-2013.




Figure 2 Unit hydrograph of the Zahedan aquifer in the period 2002-2014

The Gaussian model for parameters of  $K^+$ ,  $SO_4^{2-}$ , SAR, and the sum of cations in both periods was chosen as the best model for evaluation, while the spherical model for parameters of  $Na^+$ ,  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Cl^-$ , and TDS was recognized as the best model. For the parameters of  $HCO_3^-$ , pH, and EC and the sum of anions in 2002-2003 the spherical model and in 2012-2013 the Gaussian model provided higher estimation accuracy.

### 3.5 Zoning of the chemical parameters of ground water quality

Considering the large number of maps prepared for different years, the first period (2002-2003) was chosen as the sample of dry year, while the second period (2012-2013) was selected as the normal year sample for presenting zoning maps.

#### 3.5.1 Electrical conductivity (EC)

Figure 3 demonstrates zoning maps of electrical conductivity for the periods of 2002-2003 and 2012-2013. EC in 2012-2013 had increased in the northeast, eastern, and southeastern parts of the aquifer.




Figure 3 Changes in EC of the Zahedan aquifer in the dry and normal periods

#### 3.5.2 Total dissolved solids (TDS)

Comparing the two maps, TDS levels had increased in the northern, northeastern, and

eastern parts of the aquifer during the period 2012-2013 (Figure 4).




Figure 4 Changes in total dissolved solids of the Zahedan aquifer in the dry and normal periods

### 3.5.3 Acidity (pH)

The ground water pH had become more alkaline during the period 2002-2003 in the

central and southern parts of the Zahedan Plain's aquifer (Figure 5).




Figure 5 Changes in acidity of the Zahedan aquifer in the dry and normal periods

### 3.5.4 The sum of actions

The sum of cations in the normal period (1391-1392) had increased in the northeastern,

western, and southern parts of the aquifer (Figure 6).




Figure 6 Changes in total cations of the Zahedan aquifer in the dry and normal periods

### 3.5.5 The sum of anions

The sum of anions in the normal period (1391-1392) had increased in the northern, eastern,

and the southern parts of the aquifer, while it had declined in the northeastern, western, southern, and central parts (Figure 6).




Figure 7 Changes in total anions of the Zahedan aquifer in the dry and normal periods

### 3.6 Classification of groundwater for consumption

#### 3.6.1 Water quality in terms of drinking

Using Schoeller diagram, all the wells were chemically classified for drinking within both

normal and dry periods (Figures 8 and 9), which showed that the quality of potable water in Zahedan Plain had not changed significantly within the dry and normal periods, thus suitable for drinking.




Figure 8 Classification of the Zahedan plain wells for drinking in the dry period





Figure 9 Classification of the Zahedan plain wells for drinking in the normal period




### 3.6.2 Quality in Terms of Agriculture

Wilcox classification for agricultural use indicated that the ground water quality of the Zahedan Plain within the dry and normal

periods varied between the limits of  $C_3S_1$  and  $C_4S_1$  (Figures 10 and 11). This suggests that drought had not had significantly affected the quality of the water for agricultural purposes.



**Figure 10** Classification of the Zahedan plain wells in the dry period for agricultural use



**Figure 11** Classification of the Zahedan plain wells in the normal period for agricultural use

### 3.6.3 The Piper Diagram

Based on the Piper Diagram, the ground waters of the Zahedan Plain within the dry period had three types of sodic chloride, calcic

chlorine, and sodic sulfate (Figure 12), while two types of sodic chloride and sodic bicarbonate were recognized within the normal period (Figure 13).




Figure 12 Piper diagram of the Zahedan plain wells in the dry period




Figure 13 Piper diagram of the Zahedan plain wells in the normal period

### 3.7 Correlation between the quality parameters and the balance of ground water level with drought indices

Table 3 lists the correlation between the water quality parameters and drought indices of SPI and PN. A significant correlation between the sum of anions and TDS with SPI was observed

at 5% level. There was also a significant correlation between  $\text{Na}^+$  and SPI at 1% level. However, there was no significant correlation between the quality parameters and PN. A significant correlation at 1% level between the water balance and only SPI at the level of 1% was found.

**Table 3** Correlations between quality and ground water level in Zahedan plain with drought indices SPI and PN

		EC	TDS	pH	CL <sup>-</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	Na <sup>2+</sup>	Cation	SO <sub>4</sub> <sup>2-</sup>	HCO <sub>3</sub> <sup>-</sup>	Anion	ground water level (m)
SPI	Pearson Correlation	.421	.637*	-.143	.451	.501	.574	.412	.741**	.599	.453	.144	.637*	-.869**
	Sig. (2-tailed)	.197	.035	.674	.164	.117	.065	.208	.009	.051	.162	.672	.035	.001
PN	Pearson Correlation	-.119	-.006	.122	-.268	.141	.192	-.217	-.210	-.126	-.087	.322	-.155	-.043
	Sig. (2-tailed)	.727	.987	.720	.425	.679	.572	.521	.536	.711	.800	.334	.649	.901

\*\* The parameters significantly correlated with the level of 1%

## 4 DISCUSSION AND CONCLUSION

In many studies, only SPI has been used for investigation of ground water, but in this research the effect of PN index on the ground water resources was also taken into consideration along with SPI. The results of the drought index of PN indicated that this index provided a greater variety than SPI. However, for drought studies and its impact on the ground water resources, it does not have the necessary efficiency. The results of the ground water level and unit hydrograph of the aquifer indicated a 1.37 m elevation in the phreatic zone in Zahedan aquifer from the periods of 2002-2003 to 2013-2014 (Fig. 2), which is in contrast with studies in other regions (e.g. Watmough and Juckers, 2014; Najafzadeh *et al.*, 2015). Considering the drought conditions, this increase in the ground water level is because that the water requirements of citizens in Zahedan city (over 600 thousand people requiring water about 25 million cubic meters in the year) has been supplied from another source called Chah-Nimehs (fed by the Hirmand River), which in turn decreased

exploitation of water from the wells across the aquifer. Moreover, lack of waste water collection system in Zahedan City and seepage of the disposed water through the plain's aquifer has also contributed to increased level of the ground water (Sarhadi, 2015; Regional Water Company, 2011). Currently, 95% of the urban population in Zahedan is under coverage of water distribution system.

The findings of this section of the research are in harmony with the results obtained by Ahmadvash *et al.* (2013) considering elevated ground water level. There was a correlation between water level and the SPI drought index at 1% level. Investigation of the trend of changes in EC, dry residual, and the sum of cations and anions of this aquifer (Figs. 3 to 7) demonstrated that the extent of these parameters were minimal in the west of the aquifer. It then dramatically grows at the direction of ground water movement towards east and north, which can be attributed to the presence of evaporative and quality-lowering sediments in the aquifer. The results of correlation of the water quality parameters

with SPI indicated that there was a significant correlation at 1% level between  $\text{Na}^+$  and SPI, but no significant difference was observed between the quality parameters and PN (Table 3). Regarding correlation between the ground water level and the quality parameters with SPI drought index, the findings of this research are in correspondence with those obtained by Aleboali *et al.* (2016), Jahanshahi *et al.* (2016) and Khan *et al.* (2008). Determination of the consumability of the ground waters of Zahedan Plain using Schoeller and Wilcox diagrams showed that drought had no significant effect on the class of ground water quality for drinking and agricultural purposes. Finally, considering the significance of water transfer project from Sistan Chah-nimehs to Zahedan, it is suggested that the environmental and socio-economic effects of this project should be investigated on water and soil resources in the area of origin (Sistan) and destination (Zahedan). Since farmers and other exploiters highly rely on the ground water resources for their requirements, permeation of salty waters towards the plain and its consequences should be considered and the necessary measures to preserve the plain's water resources should be investigated.

## 5 REFERENCES

- Afzali, A. and Shahedi, K. The trend of changes in ground water quality and quantity of Babol- Amol. *Journal of Watershed Management \ Research*. 2014; 5(1): 144-156. (In Persian)
- Ahmadvash, S., Babazadeh, H., Kaveh, F. and Bybordi, A. The effects of the recent drought in terms of quantity and quality of ground water resources in Azarshahr. The first national conference on the water crisis, Islamic Azad University, Khorasgan (Esfahan), Iran, 25-26 May 2013. 11 p. (In Persian)
- Ahmed, S. Ground water monitoring network design: Application of geostatistics with a few case studies from a granitic aquifer in a semi-arid region, in ground water hydrology, M.M. Sherif, V.P. Singh and M. Al-Rashed (Eds.), Balkema, Tokyo, Japan, 2002; 2: 37-57.
- Akbari, M., Jargeh, M.R. and Sadat, M.H. Assessment of decreasing ground water level using geographic information system (GIS) (case study: Mashhad plain aquifer). *Journal of Soil and Water Conservation*. 2009; 16(4): 78-63. (In Persian)
- Aleboali, A., Ghazavi, R. and Sadatinezhad, J. Study the effects of drought on ground water resources using spi index (a case study: Kashan plain). *Desert Ecosystem Engineering Journal*. 2016; Issue 10, 13-22. (In Persian)
- Arsalan, H. Spatial and temporal mapping of ground water salinity using ordinary kriging and indicator kriging: the case of bafra plain, Turkey. *Agr. Water Manage.*, 2012; 113: 57-63.
- Goovaerts, P. Geostatistics for natural resources evaluation. Oxford University Press, New York, 1997; 483 P.
- Jahanshahi, A., Moghddam Nia, A.R. and Nohtani, M. Assessment of drought impact on ground water resources using SPI index (case study: Shahr-e-Babak plain, Kerman province). *Desert Ecosystem Engineering Journal*. 2016; Issue 10, 85-98. (In Persian)
- Jahanshahi, A., Rouhi Moghadam, E. and Dehviri, A. Assessment of ground water quality parameters using gis and geostatistical (case study: shahr-e-babak plain). *Journal of Soil and Water Science*, 2014; 24(2): 183-197. (In Persian)

- Juckers, M. and Watmough, S.H.A. Impacts of simulated drought on pore water chemistry of peatlands. *Environ. Pollut.*, 2014; 184: 73-80.
- Khan, S., Gabriel, H.F. and Rana, T. Standard precipitation index to track drought and assess impact of rainfall on water tables in irrigation areas. *J. Irr. Drain. Syst.*, 2008; 22: 159-177.
- McKee, T.B., Doesken, N.J. and Kleist, J. The relationship of drought frequency and duration to time scales, *Proceedings of the English Conference on Applied Climatology*, Boston, MA: American Meteorology Society, 1993; 179-184.
- Nahvinia, M.G., Nikmehr, C., Aly, Kh.A., Ramazani, H. and Kolahchi, Aa. Spatial analysis of drought indices SPI and SIAP in the province Kerman, *First International Conference on Water Crisis*, Zabol University, Iran, 20-22 March 2008. (In Persian)
- Najafzadeh, H., Zehtabian, Gh.R., Khosravi, H. and Golkarian, A. Climatic and geological factors impact on the quantity and quality of ground water resources mah-velayat plain. *Ecohydrology*. 2015; Volume 2, Issue 3, 325-336. (In Persian)
- Nosrati, K. and Eeckhaut, M.V.D. Assessment of ground water quality using multivariate statistical techniques in Hashtgerd plain, Iran. *Environmental Earth Science*, 2011; 65: 331-344. (In Persian)
- Regional Water Bureau. *Water resources of Zahedan*, 2002. (In Persian)
- Sarhadi, M. The effect of drought on ground water quantity and quality of zahedan plain, *Watershed Master's Thesis*, University of Zabol, 2015; 137 P. (In Persian)
- Schoeller, I. *Les eaux souterraines, massoneteic*. 1962; 923 p.
- Taghizadeh Mehrjerdi, R., Zareian, M., Mahmodi, Sh. and Heidari, A. Spatial distribution of groundwater quality with geostatistics (case study: Yazd-Ardakan plain). *World Applied Sciences Journal*, 2008; 4(1): 7-9.
- Wilcox, L.V. *Classification and use of irrigation water*. U. S. Department of Agriculture, Washington, 1955; 969 P.

## اثر خشکسالی بر پارامترهای کمی و کیفی آبخوان دشت زاهدان

معصومه سرحدی<sup>۱</sup>، محمد نهتانی<sup>۲\*</sup> و محمد ریگی<sup>۳</sup>

- ۱- دانش آموخته کارشناسی ارشد، گروه مرتع و آبخیزداری، دانشگاه زابل، زابل، ایران
- ۲- استادیار گروه مرتع و آبخیزداری، دانشکده آب و خاک، دانشگاه زابل، زابل، ایران
- ۳- دانش آموخته کارشناسی ارشد، گروه مرتع و آبخیزداری، دانشگاه زابل، کارشناس ظرفیت‌سازی پروژه بین‌المللی مناخ در استان سیستان و بلوچستان، ایران

تاریخ دریافت: ۱۳ اردیبهشت ۱۳۹۵ / تاریخ پذیرش: ۲۵ مرداد ۱۳۹۵ / تاریخ چاپ: ۱۹ آذر ۱۳۹۵

چکیده با استفاده از شاخص‌های خشکسالی درصد نرمال بارش (PN) و بارش استاندارد (SPI)، تاثیر خشکسالی بر تغییرات سطح ایستابی و پارامترهای کیفی آبخوان دشت زاهدان بررسی شد. برای بررسی تغییرات کیفیت و سطح ایستابی آبخوان از آمار ۱۲ ساله چاه‌های مشاهده‌ای و پی‌زومتری در دشت و برای بررسی شاخص‌های خشکسالی PN و SPI، از آمار ۳۰ ساله بارندگی ایستگاه زاهدان استفاده شد. نتایج بیانگر تنوع بیش‌تر شاخص PN نسبت به شاخص SPI بود. هیدروگراف معرف آبخوان نیز بیانگر افزایش ۱/۳۷ متری سطح ایستابی از سال آبی ۲۰۰۲-۲۰۰۳ تا ۲۰۱۳-۲۰۱۴ بود که با شرایط خشکسالی می‌تواند به علت کاهش برداشت از آبخوان در پی انتقال آب از چاه‌نیمه‌های سیستان به زاهدان باشد. همبستگی معنی‌داری بین شاخص SPI و پارامترهای کیفی آب همچون TDS، Anion و Na<sup>+</sup> با SPI مشاهده شد، اما شاخص PN با هیچ‌کدام از پارامترهای کیفیت همبستگی معنی‌داری نداشت. در مورد تراز سطح آب نیز فقط با شاخص SPI همبستگی معنی‌داری در سطح ۱ درصد مشاهده گردید.

**کلمات کلیدی:** دشت زاهدان، شاخص‌های خشکسالی، کیفیت آب زیرزمینی، هیدروگراف واحد آبخوان