

Effect of sowbug on Soil Aggregate Stability in a Desert Region (Case Study: Gareh Bygone Plain, Iran)

Gholamreza Rahbar^{1*}, Ataollah Kavian², Mahmood Habibnezhad Rooshan³, Ahang Kowsar⁴ and Kaka Shahedi²

¹ Ph.D. Candidate, Department of Watershed Management, Faculty of Natural Resources, Sari Agricultural Science and Natural Resources University, Sari, Iran

² Associate Professor, Faculty of Natural Resources, Sari Agricultural Science and Natural Resources University, Sari, Iran

³ Professor, Faculty of Natural Resources, Sari Agricultural Science and Natural Resources, Sari, Iran

⁴ Professor, Department of Soil Conservation and Watershed Management, Fars Research Centre for Agriculture and Natural Resources University, Shiraz, Iran

Received: 18 October 2015 / Accepted: 21 December 2015 / Published Online: 9 January 2016

ABSTRACT The appearance of sowbugs (*Hemilepistus shirazi* Schuttz) in the sedimentation basins of the artificial recharge of groundwater (ARG) systems in the Gareh Bygone Plain (GBP) in southern Iran is considered an ecological breakthrough in desertification control. This study was performed at the Kowsar Floodwater Spreading and Aquifer Management Research, Training and Extension Station in GBP, 200 km from the south east of Shiraz, Iran, on the alluvial fan of Bisheh Zard River. Invasion of sowbug to sedimentation of basin due to water increasing persuade us to study about this crustacean. To determine aggregate size distribution, the soil samples were dried, and then the soil was sieved through a set of sieves (8, 4.75, 2.8, 2.0, 1.0, 0.8, 0.3 and <0.3 mm) and the aggregate size distribution was determined. Results also showed that the mean aggregate stability percentage for control (without sowbug) was 26.14% and for burrowed materials (with sowbug) was 78.00%. The aggregate stability percentage of burrowed materials was 3 folds more than that of control. The mean weight diameter for burrowed materials of sowbug was 1.86 g against 0.44g for control. Domestication of this useful organism, a souvenir of the ARG systems is an environmentally sound and a financially viable method of lengthening the economic life of the artificial recharge of groundwater system.

Key words: Desert management, Floodwater spreading, Soil stability, Terrestrial isopod

1 INTRODUCTION

Groundwater consists about 70% of the water consumed in Iran during periods with normal precipitation and a higher percentage during droughts (Kowsar, 2013). Over-exploitation of groundwater in recent droughts has left our agricultural sector in the imminent danger of

collapse. However, desertification control through floodwater spreading, particularly for the artificial recharge of groundwater (ARG) on a very large scale, may delay this ultimate crisis, mainly due to reclamation of the drastically disturbed land and recharging the badly depleted aquifers. Implementation of

* Corresponding Author: Ph.D. Candidate, Department of Watershed Management, Faculty of Natural Resources, Sari Agricultural Science and Natural Resources University, Sari, Iran, Tel:+98 917 708 1360, E-mail: gholamreza.rahbar@gmail.com

the ARG causes sedimentation of the nutritious suspended load, which are abundant in floodwater, thus soil building in the deserts underlain with potential aquifers (Kowsar, 2005; Kowsar, 2011).

It is well-known that various macro fauna inhabit the soil and modify some of its properties (Pankhurst and Lynch, 1994). Tunneling, mounding and pedoturbation are some of the most observable activities which have attracted the attention of soil scientists. Ants, beetles, termites, cicadas and crayfish are a few of the fauna whose activities, which somehow resemble those of the sowbug, have been studied (Tiedeman, 2013). Measurements of aggregate stability have received considerable attention during the last 50 years. It is generally agreed that soil structure stability depends mainly on forces that bind microaggregates (Tisdall and Oades, 1982) into macroaggregate. Macroaggregation depends primarily on temporary binding agents and is believed to be sensitive to changes in soil organic matter content induced by soil management (Williams, 1971; Channey and Swift, 1986; Besalatpour *et al.*, 2013).

Earthworms, known to increase aggregate size and stability (Ziegler and Zech, 1992). In recent years, Margonelli (2014) studied termite mounds and its structure carefully and published it in National Geographic in August 2014. The latter investigator discovered indicator plants for the geological formations where specific species of termites were active. Luken and Kalisz (1984) reported that three cicada species (*Magicicada cassini*, *M. septendecim*, and *M. septendecula*) that live underground for a period of either 13 or 17 years, burrow to a depth of 36 cm. Stone (1993) found that a crayfish (*Procambarus rogersi* Hobbs.) excavates galleries four to 10 cm in diameter, and longer than 1.5 m, chiefly in the upper 30 cm of soil, but with vertical shafts deeper than one m. Ketterings *et al.* (1997) investigated the impacts of earthworms on soil

aggregate stability in a legume cover crop agro-ecosystem. They suggested that earthworm's activity can increase soil structure stability. Although all of the burrows made by the above mentioned animals could function as flow path for water and solutes, this point was not an objective of these investigations. An aggregate is a group of primary particles that cohere to each other more strongly than to other surrounding soil particles. Aggregate stability on the other hand is measure of aggregate resisting disruptive force (Kemper and Rosenau, 1986). Aggregate size distribution after dry sieving can be expressed as a single empirical unit called mean weight diameter (MWD) which is an important characteristic because, firstly, resistance to wind and water erosion and secondly, it give an idea of the distribution of the large pores after tillage. Aggregate stability of soil is a key factor of soil resistivity to mechanical stresses, such as the impacts of rainfall and surface runoff, and therefore to water and wind erosion, which might be prevented by mulching the soil surface. The measurement of soil aggregate stability becomes important, because it can give general information about soil condition (Canasveras *et al.*, 2010). The aggregate stability is the ability of the bonds of the aggregates to resist when exposed to stresses causing their disintegration. Information on soil aggregate stability can also improve programs, adapted to the specific soil type and crop demands (Chenu and Plante, 2006).

The objective of this study is to evaluate the role of sowbug in the formation of aggregates of different sizes and soil aggregate stability changes in the floodwater spreading systems in Gareh Bygone Plain in Fars Province.

2 DESCRIPTION OF THE SOWBUG

Sowbug is a crustacean, 10-15 mm long and five mm wide. It is blackish gray and has seven pairs of legs (Figure 1). According to last classification of crustacean (Barnes, 1987.

Martin and Davis, 2001) and terrestrial Isopods (Schmidt, 2003), this organism has been classified as follows:

Category Metazoa

Phylum Arthropoda

Subphylum Crustacea

Class Malacostraca Latreille, 1806

Order Isopoda Latreille 1803

Suborder Oniscidae Latreille 1803

Genus *Hemilepistus*

Species *Hemilepistus shirazi* Schuttz

sowbugs live in damp places, forage on vegetation, and digest the soil organic matter. Their burrows, seven mm in diameter and up to 185 cm deep, serve to aerate and drain the soil profile. The same genus or one of its close relatives burrows down to 100 cm in Gonabad, Northeast of Iran (Rahimi, 1993).

A species of *Hemilepistus* has been shown to excavate up to 1.5 t ha^{-1} in its active period of about three months in Central Asia. The excavated soil had more organic matter and a better structure, thus was more resistant to

erosion than the original soil from which it was extracted (Salehrastin, 1978).

This viviparous organism lives for about one year. The white brood pouch under the abdomen of the female swells in March. The eggs form larvae in the pouch, and 60-70 sowbugs, very similar to their parents, are released from the pouch in May. They are very active in the spring and fall. They come out of their burrows in the cool air of early morning and late afternoon. It seems that digging deep into the soil is to reach a humid surrounding. There are semi-spherical spaces at the end of their burrows, five to 10 cm in diameter (Rahbar *et al.*, 2015). They form semi-cylindrical, rods of soil, two mm long and one mm in diameter (burrowed materials), with their mandibles, and place them to one side of the opening of their burrows (Figure 2). These rods are seemingly more resistant to slaking than the freshly laid sediment from which they are formed. As some remains of the sowbug are found in scorpion burrows, it is believed that *Hemilepistus shirazi* Schuttz is eaten by this arachnid.



Figure 1 A dorsal view of a female *Hemilepistus shirazi* Schuttz (Rahbar *et al.*, 2015)



Figure 2 Parallelepiped-shaped burrowed materials of sowbug

3 MATERIALS AND METHODS

3.1 Study area description

The study was performed at the Kowsar Floodwater Spreading and Aquifer Management Research, Training and Extension Station, and lies between the latitudes 28°35` and 28°40` N and the longitudes 53°53` and 53°57` E on debris cone in the Gareh Bygone Plain (GBP) that is a small island of drifting sands and moving dunes of Iran (Kowsar, 1991). The field study is about 200 km to the Southeast of Shiraz, Iran (Figure 3). The soil

texture of the site is loamy sand [coarse-loamy over loamy skeletal, carbonatic, (hyper) thermic, Typic Haplocalcids]. Eight ARG systems, covering a total area of 1365 ha, were installed during the 1983-1988 period (Kowsar, 1998., Kowsar, 2005).

The walls of sowbug burrows are quite durable. Apparently, the sowbug wet and cement their burrows with the body fluids. An extremely thin coating of a grayish material lines the burrows in very fine sand that collapse otherwise.




Figure 3 The location of study area in Fars Province, Iran

3.2 Experiment design

One of the floodwater spreading systems that have six plots was selected (Figure 4). Twenty four samples (2 kg) of burrowed materials were taken from near the burrows of sowbug which is about 250 ha of the site. As control, about 2 kg of the rod shaped soil surface were taken from adjusted plots without the burrows and sowbugs. After air dry of soil samples sent them to soil physical laboratory of Fars research center for agriculture and natural resources. The most common method used for aggregate stability measurement is wet sieving. Aggregate stability measured using a wet-sieving

apparatus (Eijklkamp, Netherlands). The method of wet-sieving is adapted from Kemper and Rosenau (1986). Aggregate stability is expressed as the percentage of aggregates remaining larger than 0.25 mm. Correction for sand particles larger than 0.25 mm must be than as these particles cannot be broken down further to pass through the sieve (Nimmo and Perkins, 2002; Teh and Jamal, 2006).

Aggregate stability of soil samples was computed by Eq. 1 (Teh and Talib, 2006).

$$AS\% = \frac{W-S}{4-S} \times 100 \quad (1)$$



Figure 4 Location of the 2nd sedimentation basin of Bisheh Zard 1

Where AS is aggregate stability, Wis the weight of oven dried- aggregates and S is the weight of sand particle.

To determine aggregate size distribution and mean weight diameter (MWD) the soil samples were dried, and then 1500 g of the soils were passed through a set of sieves (8, 4.75, 2.8, 2.0, 1.0, 0.8, 0.3, and <0.3 mm).

The weight of aggregates remaining on each sieve was used to determine aggregate size distribution. Aggregate size distribution after dry sieving can be expressed as an empirical unit called MWD. This was calculated using the Eq. 2 (Teh and Talib, 2006).

$$MWD = \sum_{i=1}^n Xi Wi(2)$$

where MWD is mean weight diameter, Xi is the mean diameter and Wi is the proportion of the total sample weight.

3.3 Statistical analyses

Statistical analyses were performed using SAS and SPSS 19 softwares. At first normality of data verified through Kolmogrov and Smirnov test and data homogeneity of variance was

evaluated by Leven test. Analysis of variance (ANOVA) in a completely randomized design was used to determine the effect of treatments for each measured variable. As there were only two treatments, the burrowed materials and the freshly-laid sediment (control) the paired t-test was used to detect the significance effect of the sowbug activities on study variables ($P < 0.01$).

4 RESULTS AND DISCUSSION

The results of aggregate size distribution and aggregate stability of burrowed materials and soil control are shown in Tables 1 and 2 and Figure 5. Analysis variance results on soil aggregate stability of sowbug burrowed materials and soil samples (control, without sowbug activity) shows that there was significantly higher ($P < 0.01$). Table 1 and Figure 5 illustrate the mean aggregate stability percentage for study treatments. The mean aggregate stability of control treatment (without sowbug) was 26.14% and of burrowed materials (with sowbug) was 8.00%. The aggregate stability percentage of burrowed materials was threefolds more than the control.

In Table 2, the MWD for burrowed materials of sowbug was obtained 1.86 g and 0.44 g for the control and it was more than four folds in comparison with control.

Larger values of MWD denote a higher proportion of larger aggregates in soil sample. The index of MWD is developed to give greater weighting to larger aggregates than that for smaller aggregates. This reflects the fact that larger aggregates often indicate favorable soil structure for agriculture. Aggregate size distribution can also be expressed as the

percentage of aggregates that are greater than 2 mm. Soil aggregation is a fundamental property of soils and is a primary control of soil aeration. It determines the hydrological properties of soil such as water-holding capacity and the storage of organic carbon. The stability of soil aggregates is also important because it influences how these properties change with time, and the susceptibility of soils to erosion by both wind and water which can lead to a loss of soil nutrients (Rawlins *et al.*, 2013).

Table 1 Comparison between the control and burrowed materials

Group	N	Means	Significant level
Control (without sowbug)	8	26.14 ± 4.19 ^b	<0.01
Burrowed materials (with sowbug)	24	78.00 ± 5.56 ^a	




Figure 5 Aggregate stability percentage for control and burrowed materials (BM)

Table 2 The mean weight diameter (MWD) of control and burrowed materials (BM) treatments

Size (mm)	8.0	4.75	2.80	2.0	1.0	0.80	0.30	<0.30	MWD
BM Weight (g)	0	0.30	5.53	95.0	145.5	4.75	1.89	2.0	1.86
Control Weight (g)	0	0	0	0.23	6.63	49.82	106	98.92	0.49

Some large soil invertebrates have significant effects on soil structural properties, the most important being earthworms, termites and ants. They built organo-mineral structure of different stability such as galleries, casts, fungous-comb chambers and mound (Lavell *et al.*, 1997). Activity of the sowbug is beneficial for improvement of soil physical conditions and thus, plant growth. Burrowing by the sowbug improves air and water penetration into the soil. The burrowed materials and castings resist erosion more than the freshly-laid sediment from which they are formed (Jass and Klausmeier, 2000; Schmidt and Leistikow, 2005). The mechanisms by which sowbug increased soil aggregate stability are not well understood. Much of the research to date has focused on casts. Burrow materials stability is dependent in part on the quality of the ingested organic matter; quality of C (carbon) in the casts may be a factor affecting their stability. Farid, *et al.* (2014) state that organic matter has the highest direct and positive effect on soil aggregate stability in the north west of Iran and it causes increasing stability of soil aggregate. Soil organic matter and texture (clay content) are the main a biotic binding agents in the formation and stabilization of aggregates and play key roles in controlling soil structure stability through their influence on water absorptivity and repellency as well as on the strength of bonds between particles, while soil microbes (bacteria and fungi) and plant roots have been reported as key biotic aggregating agents (Duchicela *et al.*, 2013). Because sowbug eat residue of plants they can increase the contents soil organic carbon and their role can be similar to earthworms, termites and ants.

5 CONCLUSION

sowbugs are incredible organisms that sustain floodwater spreading systems in the south of Iran in Gareh Bygone Plain, and they are important biological factors in soil ecosystem.

The influence of sowbug on soil aggregate stability is very important for maintenance of soil fertility and quality. The results of this research stated that sowbugs had significant and positive effects on soil aggregate stability, and showed that this terrestrial organism can make a soil with better quality. Nevertheless further research regarding the impact of sowbug on soil quality is required. In addition, it will be important to explore the effects of sowbug in chemical properties of soil.

6 REFERENCES

- Barnes, D.R. Invertebrate zoology. Philadelphia, pa: holt-saunders international. 1987; 743 P.
- Besalatpour, A.A., Ayobi, S., Hajabbasi, M.A., Mosaddeghi, M.R. and Rainer S. Estimating wet soil aggregate stability from easily available properties in a highly mountainous watershed. *Catena*, 2013; 111:72-79.
- Chaney, K. and Swift, R.S. Studies on aggregate stability. 1- reformation of soil aggregate. *J. Soil Sci.*, 1986; 37:329-335.
- Chenu, C. and Plante, A.F. Organic matter influence on clay wettability and soil aggregate stability. *Soil Sci. Soc. Am. J.*, 2006; 64: 1479-1486.
- Conasveras, J.C., Barron, V., Campillo, M.C., Torrent, J. and Gomez, J.A. Estimation of aggregate stability indices in Mediterranean soil by diffuse reflection spectroscopy. *Geoderma.*, 2010; 158: 78-84.
- Duchicela, J., Sullivan, T., Bontti, E. and Bever, J. Soil aggregate stability increase is strongly related to fungal community succession along an abandoned agricultural field chronosequence in the

- Bolivian Altiplano. *J. Appl. Ecol.*, 2013; 50: 1266-1273.
- Farid, B., Arami A. and Akhsari, D. Assessing the role of some soil properties on aggregate stability using path analysis (Case study: silty clay-loam and clay loam soil from Gully land in North West of Iran). *ECOPERSIA.*, 2014; 2(2): 513-523.
- Jass, J. and Klausmeier, B. Endemics and immigrants: North American terrestrial isopods (Isopoda, Oniscidea) north of Mexico. *J. Crustacean. Biol.*, 2000; 73: 771-799.
- Kemper, W.D. and Rosenau, R.C. Aggregate stability and size distribution. In methods of soil analysis. Part 1. Physical and mineralogical methods, 2nd ed. ASA-SSSA, Wisconsin, 1986; 425-442.
- Kettering, Q.M., Blair, J.M. and Marinssen, J.C.Y. Effects of earthworms on soil aggregate stability and Carbon and nitrogen storage in legume cover crop agroecosystem. *Soil Biol. Biochem.*, 1997; 29(3/4): 401-408.
- Kowsar, A. Abkhandari (Aquifer Management): a green path to the sustainable development of marginal drylands. *J. Mont. Sci.*, 2005; 2(3): 233-243.
- Kowsar, A. Desertification control through floodwater harvesting: the current state of know-how. The future of drylands, Chapter Five UNESCO. 2011; 229-241.
- Kowsar, A., Rahbar, Gh., Vali, A., Nejabat, M., Mohammadnia, M., and Jowkar, L. An evaluation of geological nitrogen in the Roudshur-Jahrom, Watershed (Final report of research plan). *Fars Res. Cent. Agr. Nat. Res.*, 2013; 65 P.
- Kowsar, S.A. Aquifer management: A key to food security in the deserts of the Islamic Republic of Iran. *Desertification Control Bulletin (UNEP).*, 1998; 33:24-28.
- Kowsar, S.A. Floodwater spreading for desertification control: An integrated approach. *Desertification Control Bulletin (UNEP).*, 1991;19: 3-18.
- Lavell, P., Bignell, D., Lepage, M., Volters, V. and Rogers, P. Soil function in a changing world: the role of invertebrate ecosystem engineers. *Eur. J. Soil Biol.*, 1997; 33:159-193.
- Luken, J.O. and Kalisz, J. Soil disturbance by the emergence of periodical cicadas. *Soil Sci. Soc. Am. J.*, 1989; 53 (1): 310-313.
- Margonelli, L. Collective mind in the mound: How do termites build their huge structure? *Natl. Geogr.*, August 2014.
- Martin, J.W. and Davis, G.E. An updated classification of the recent crustacea (Science series / Natural History Museum of Los Angeles County). Publisher: Natural History Museum of Los Angeles County, 2001; 388 P.
- Nimmo, J.R. and Perkins, K.S. Aggregate stability and size distribution. Methods of soil analysis. Part 4. Madison, Wisconsin, Soil Soc. Am., 2002; 314-328.
- Pankhurst, C.E. and Lynch, J.M. The role of soil biota in sustainable agriculture. *Soil biota: Management in sustainable farming systems.* CSIRO, Australia. 1994; 3-8.
- Rahbar, Gh., Kowsar, A. and Azimi, M. Bioassay of sowbugs, burrowed material using barley and wildrye. Second National Conference on Desert, Semnan, Iran. 2015. (In Persian)

- Rahimi, H. Morphological and biological characteristics of sowbug. *Zeitun: Scientific and Specific Monthly in Agriculture*, 1993; 115: 36-38. (In Persian)
- Rawlins, B.G., Wragg, J., Lark, M. Application of a novel method for soil aggregate stability measurement by laser granulometry with sonication. *Eur. J. Soil Sci.*, 2013; 64(1): 92-103.
- Salehrastin, N. Soil biology: (Soil inhibiting organisms and their role in distribution of elements). Tehran University Publication, 1978; 1666: 482 P. (In Persian)
- Schmidt, C. and A. Leistikow. Review of the genus *Androdeloscia* Leistikow, with description of four new species (Crustacea: Isopoda: Oniscidea). *Entomologische Abhandlungen*, 2005; 62 (2): 117-163.
- Schmidt, C. Contribution to the phylogenetic system of the Crinocheta (Crustacea, Isopoda). Part 2 (Oniscidae to Armadillidiidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin (Zoologische Reihe)*, 2003; 79: 3-179.
- Stone, E.L. Soil burrowing and mixing by a crayfish. *Soil Sci. Soc. Am. J.*, 1993; 57:1096-1099.
- Teh, C.B.S. and Talib, J. Soil physics analysis. University Putra Malaysia Press. 2006; 42 P.
- Tiedeman, M.L. Pedogenesis of ant-colonized soil at Doolittle Parairie State Preserve, Iowa. M.Sc. Thesis, Iowa State University. 2013. 63 P.
- Tisdall, J. M. and Oades, J.M. Organic matter and water-stable aggregate in soil. *J. Soil Sci.*, 1982; 33:141-163.
- Williams, R.J.B. Relationship between composition of soils and physical measurements made on them. 1971; Rthamsted Experimental Station. Report for 1971- Part 2.
- Ziegler, F. and Zech, W. Formation of water-stable aggregates through the action of earthworms. *Pedologia.*, 1992; 36:91-96.

تأثیر خرخاکی بر پایداری خاک دانه‌های خاک یک منطقه بیابانی (مطالعه موردی: دشت گربایگان فسا، ایران)

غلامرضا رهبر^{۱*}، عطاء ا. کاویان^۲، محمود حبیب‌نژاد روشن^۳، آهنگ کوثر^۴ و کاکا شاهدی^۲

- ۱- دانشجوی دکتری علوم و مهندسی آبخیزداری، دانشکده منابع طبیعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ایران
- ۲- دانشیار، دانشکده منابع طبیعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، ایران
- ۳- استاد، دانشکده منابع طبیعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، ایران
- ۴- استاد پژوهش، بخش تحقیقات حفاظت خاک و آبخیزداری، مرکز تحقیقات کشاورزی و منابع طبیعی فارس، شیراز، ایران

تاریخ دریافت: ۲۶ مهر ۱۳۹۴ / تاریخ پذیرش: ۳۰ آذر ۱۳۹۴ / تاریخ چاپ: ۱۹ دی ۱۳۹۴

چکیده حضور خرخاکی‌ها در حوضچه‌های تغذیه مصنوعی شبکه‌های پخش سیلاب گربایگان فسا در جنوب شرقی ایران به‌عنوان راهی برای مهار بیابان‌زایی در نظر گرفته می‌شود. خرخاکی یکی از موجودات خاک‌زی مفید است که سبب نفوذ بیش‌تر آب به خاک و سبب بهبود ویژگی‌های فیزیکی خاک شده است. از خاک‌هایی که از فعالیت خرخاکی‌ها به‌دست آمده و هم‌چنین نقاط شاهد (مکان‌های بدون فعالیت خرخاکی) نمونه‌هایی برداشت شد. بعد از هوا خشک شدن نمونه‌های برداشت شده پایداری خاک دانه‌ها و هم‌چنین توزیع اندازه‌ای آن‌ها برآورد شد. برای تعیین نحوه توزیع اندازه خاکدانه‌ها نیز از سری الک‌های آزمایشگاهی (۸، ۴/۷۵، ۲/۸، ۲، ۱، ۰/۸، ۰/۳ و کم‌تر از ۰/۳ میلی‌متر) استفاده شد. نتایج نشان داد متوسط پایداری خاک دانه‌های خاک برای شاهد ۲۶/۱۴ درصد و برای خاک‌های تیمار شده با موجودات خرخاکی ۷۸ درصد بود. به‌نحوی که پایداری خاک دانه‌های مربوط به خاک‌های با فعالیت خرخاکی ۳ برابر شاهد به‌دست آمد. میانگین وزنی قطر خاکدانه‌ها برای خاک‌های حاصل از فعالیت خرخاکی ۱/۸۶ گرم و برای شاهد ۰/۴۴ گرم بود. این موجود زنده سخت‌پوست به‌عنوان راه‌آورد پخش سیلاب، سازگار با محیط‌زیست و از لحاظ اقتصادی، اجتماعی قابل قبول و سبب افزایش عمر مفید طرح‌های پخش سیلاب برای تغذیه مصنوعی سفره آب‌های زیرزمینی می‌شود.

کلمات کلیدی: پایداری خاک، پخش سیلاب، جورپایان خاک‌زی، مدیریت بیابان