

Soil Quality Indices in Pure and Mixed Forest Stands of Southern Caspian Region

Yahya Kooch^{1*}, Fatemeh Rostayee² and Seyed Mohsen Hosseini³

¹Assistant Professor, Faculty of Natural Resources, Tarbiat Modares University, Mazandaran, Noor, Iran

²M.Sc. Student, Faculty of Natural Resources, Tarbiat Modares University, Mazandaran, Noor, Iran

³Professor of Forestry, Faculty of Natural Resources, Tarbiat Modares University, Mazandaran, Noor, Iran

Received: 29 March 2015 / Accepted: 3 August 2015 / Published Online: 31 October 2015

ABSTRACT The present study aimed to assess pure planted species (*i.e.*, *Alnus subcordata* L., *Populus deltoids* L., *Taxadium distichum* L. Rich) and a mixed natural forest (*i.e.*, dominated by *Quercus castaneifolia* C. A. Mey. - *Carpinus betulus* L. - *Parrotia persica* C. A. Meyer) on basis of some soil quality indices in Mazandaran Province, northern Iran. Sixteen samples per stand were taken from the top 10 cm of soil and bulk density, texture, water content, pH, EC, organic C, total N, available nutrients, earthworm biomass, microbial respiration, fine root biomass with organic C and total N of litter layer were determined. Nine criteria (*i.e.*, silt, EC, K, Ca, Mg, microbial respiration, fine root biomass, nitrogen mineralization and litter C/N) were selected according to Principal Component Analysis (PCA) as Minimum Data Set (MDS). The analytical hierarchy process (AHP) method was employed to assign the data integration in an index. The calculated overall priority based on nine criteria, showed that the *A. subcordata* forest type had higher ecological potential (0.370) compared to the other stands. Whereas, *P. deltoids* mixed natural forest and *T. distichum* with ecological potential of 0.295, 0.213 and 0.122 had next priorities, respectively. As a conclusion, the N-fixing species, *A. subcordata*, was found more efficient in improving soil quality in degraded forest regions.

Key words: Broad-leaved species, Forest Seed Centre of Khazar, Hyrcanian Forest, Needle-leaved species, Soil characteristics

1 INTRODUCTION

The Hyrcanian or northern forests of Iran stretch up to an altitude of 2800 m above sea level and encompass different forest types thanks to the 80 tree and shrub species found there (Sagheb-Talebi *et al.*, 2014). It is obvious that these forests have been under continuous degradation over the last few decades (Kooch *et al.*, 2014), and there is an urgent need to maintain the functions of this unique forest ecosystem. National forest management

officials have acknowledged this fact and have initiated action for the sustainable management of the Caspian Forests. Different management schemes have been planned for implementation, such as documenting and exhibiting the forest disturbance, and supervising and managing the region's remaining natural forest ecosystems (Poorzady and Bakhtiari, 2009). Abandoning agriculture and tree planting for commercial or restoration purposes are also two main methods of forest restoration. There have been many

*Corresponding author: Faculty of Natural Resources, Tarbiat Modares University, Noor, Mazandaran, Iran, Tel: +98 44 55 3101-3, E-mail: yahya.kooch@modares.ac.ir

plantations of endemic and exotic species in degraded forest areas which have certainly had many effects on the ecosystem, specifically on soil fertility and nutrients. Despite the existence of extensive afforested areas and the amount of time since plantation, few studies offer a critical overall examination of the development and ecological consequences of afforestation in the Hyrcanian region (Haghdoust *et al.*, 2011), especially those related to soil properties.

Soil is an important component of terrestrial ecosystems because it preserves nutrient reserves and supports many biological processes. To preserve this resource and its functions, it is necessary first of all to know the conditions and the processes occurring in it, for example, through the determination of soil quality (SQ) (Carter, 2002; Marzaioli *et al.*, 2010). The SQ indices have been defined as soil processes and properties that are sensitive to changes in soil functions (Aparicio and Costa, 2007). It is important to build a simple, sensitive, and workable indicator method for SQ evaluation (Dumanski and Pieri, 2000). The SQ may be affected by human management practices (*e.g.* forest plantation) because these may cause alterations in soil physical, chemical and biological properties (Caravaca *et al.*, 2003).

Different methods have been developed for SQ evaluation, from qualitative or semi quantitative visual approaches (Ball *et al.*, 2007; Shepherd, 2009) to quantitative methods based on laboratory analysis and calculating SQ indices using mathematic and statistical methods (Andrews *et al.*, 2004). The SQ indices have been successfully applied at many scales and locations (Arshad and Martin, 2002; Aparicio and Costa, 2007; Mastro *et al.*, 2008). These indicators should be a combination of chemical, physical, and biological properties (Aparicio and Costa, 2007; Qi *et al.*, 2009; Lima *et al.*, 2013). Several authors have proposed sets of SQ indices (Mastro *et al.*, 2008; Marzaioli *et al.*, 2010), and have evaluated SQ based on the total data set (TDS) indicator method they selected. Also, representative indicators were

suggested by many authors, such as the minimum data set (MDS), selected according to correlation between indicators and measurement facility and the Delphi data set (DDS), selected according to the importance of the indicators to SQ based on the opinion of experts (Herrick, 2000; Zhang *et al.*, 2004; Rezaei *et al.*, 2006; Govaerts *et al.*, 2006; Zhang *et al.*, 2011).

A common feature of these based-indicator methods is that they are all identified and described by scientists and land managers according to their own terminology (Ditzler and Tugel, 2002). The present study aimed to assessment of planted species (*Alnus subcordata* L., *Populus deltoids* L., *Taxodium distichum* L. Rich) and a mixed natural forest (dominated by *Quercus castaneifolia* C. A. Mey. - *Carpinus betulus* L. - *Parrotia persica* C. A. Meyer) on basis of SQ indices in Mazandaran Province, northern Caspian region.

2 MATERIALS AND METHODS

2.1 Study area

The study area, Forest Seed Centre of Khazar, is located in the southern of Mahmudabad City, in Mazandaran Province, north of Iran. This area expands between 36° 38' N and 52° 16' E latitudes and longitudes, respectively. The study plantations composed of *A. subcordata* L., *P. deltoids* L. and *T. distichum* L. Rich. These plantations were planted in 1999 at a spacing of 4×4 m (Soleimany Rahimabady *et al.*, 2015). Beside these planted stands, a mixed natural forest dominated by *Q. castaneifolia* C. A. Mey. – *C. betulus* L. – *P. persica* C. A. Meyer was considered (Figure 1). The mean of maximum and minimum temperature were 24.4°C (in June) and 7.6°C (in December), respectively. The most of annual precipitation was 163 mm (in October) (Soleimany Rahimabady *et al.*, 2015). The climate is temperate moist and the mean of altitude from sea surface at the study site is nearly 30 m. The total slope aspect of region is facing north (Soleimany Rahimabady *et al.*, 2015).




Figure 1 Site locations of study area in Mazandaran Province, north of Iran (a) and schematic representation of the experimental design adopted for each forest stand (b)

2.2 Soil sampling and laboratory analysis

Four hectare areas (200×200 m) were selected for each stand forest in the study region. Soil sampling was carried out during the summer time using a randomly systematic method. Four soil profiles (20×20 cm) were dug along the four parallel transects in the central part of each afforested stand, thus resulting in 16 soil samples for each stand at 0-10 cm depth (Lafleur *et al.*, 2015). The same sampling procedure was carried out also for the mixed natural forest. Litter samples, simultaneously with soil samples, were collected from each stand. Total C and N contents in litter samples were determined in quadruplicate, using dry combustion with an elemental analyzer (Carrillo *et al.*, 2012). Soils were air-dried and passed through 2-mm sieve. Bulk density was measured by Plaster (1985) method (clod method). Soil texture was

determined by the Bouyoucos hydrometer method (Bouyoucos, 1962). Soil water content was measured by drying soil samples at 105° C for 24 hours. Soil pH was determined using an Orion Ionalyzer Model 901 pH meter in a 1:2.5, soil: water solution. EC (electrical conductivity) was determined using an Orion Ionalyzer Model 901 EC meter in a 1:2.5, soil: water solution. Soil organic C was determined using the Walkley - Black technique (Allison, 1975).

The total N was measured using a semi Micro - Kjeldhal technique (Bremner and Mulvaney, 1982). The available P was determined with spectrophotometer by using Olsen method (Homer and Pratt, 1961). The available K, Ca, and Mg (by ammonium acetate extraction at pH 9) were determined with Atomic absorption spectrophotometer (Bower *et al.*, 1952). The earthworms were

collected during the soil sampling by hand-sorting, washed with water and weighed. Biomass was defined as the weight of the worms after drying for 48 h on filter paper at room temperature (Jordan *et al.*, 1999). Soil microbial respiration (SMR) was determined by measuring the CO₂ evolved in 3 days incubation experiment at 25°C (Alef, 1995). Fine roots (< 2 mm diameter) were removed from each sample and dried at 70 °C to a constant mass (Neatrou *et al.*, 2005). The buried-bag technique was used to estimate soil N mineralization (Asadiyan *et al.*, 2013). Whole of these soil characters were chosen by expert opinion and literature review as important indices in soil quality (Herrick, 2000; Zhang *et al.*, 2004; Govaerts *et al.*, 2006; Rezaei *et al.*, 2006; Zhang *et al.*, 2011). Table 1 is

showing the values of studied characteristics in litter and soil layers.

2.3 Data analysis and processing

In order to synthesize all the information provided by selected parameters, a SQ index was calculated. According to Karlen *et al.* (2003), three steps are involved in the elaboration of a quality index: (1) definition of a Minimum Data Set (MDS), (2) score assignment to each indicator by mathematical functions and (3) data integration in an index. Principal component analysis (PCA) is widely used for defining a MDS and reducing data redundancy through correlation analysis among soil properties (Andrews *et al.*, 2002; Govaerts *et al.*, 2006; Li *et al.*, 2007).

Table 1 Mean values (sixteen replications in all case) of the litter and soil variable for study forest stands of *Alnus subcordata* C. A. M. (AS), *Populus deltoids* L. (PD), *Taxadium distichum* L. Rich. (TD) and mixed natural forest of *Quercus castaneifolia* C. A. Mey. - *Carpinus betulus* L. - *Parrotia persica* C. A. Meyer (QC-CB-PP)

Litter and soil features	AS	PD	TD	QC-CB-PP	F test	p value
Litter C/N	21.6 ^d	29.28 ^c	63.67 ^a	42.74 ^b	90.393	0.000
Bulk density (g cm ⁻³)	1.59 ^a	1.56 ^{ab}	1.33 ^c	1.41 ^b	5.325	0.003
Sand (%)	32.50	25.87	29.00	22.87	1.255	0.298
Silt (%)	34.62 ^b	39.75 ^{ab}	42.37 ^{ab}	46.75 ^a	3.712	0.016
Clay (%)	32.87	34.37	28.62	30.37	1.032	0.385
Water content (%)	36.76	37.75	34.56	38.72	1.165	0.330
pH (1:2.5 H ₂ O)	7.16 ^a	7.05 ^a	6.01 ^c	6.56 ^b	12.277	0.000
EC (ds m ⁻¹)	0.29 ^a	0.26 ^b	0.16 ^d	0.22 ^c	32.156	0.000
Soil C/N	4.50 ^b	5.46 ^b	28.70 ^a	10.37 ^b	11.688	0.000
Available P (mg kg ⁻¹)	24.57 ^a	22.30 ^a	12.65 ^b	15.79 ^b	15.904	0.000
Available K (mg kg ⁻¹)	337.37 ^a	328.25 ^a	156.68 ^c	257.81 ^b	35.084	0.000
Available Ca (mg kg ⁻¹)	256.25 ^a	208.68 ^b	122.81 ^d	177.31 ^c	50.674	0.000
Available Mg (mg kg ⁻¹)	57.43 ^a	52.12 ^b	36.81 ^d	44.00 ^c	34.335	0.000
Earthworm biomass (mg m ⁻²)	41.98 ^a	34.50 ^{ab}	6.23 ^c	26.26 ^b	10.696	0.000
Soil microbial respiration (mg CO ₂ -C g soil ⁻¹ day ⁻¹)	0.48 ^a	0.43 ^b	0.30 ^d	0.36 ^c	22.856	0.000
Fine root biomass (g m ⁻²)	64.60 ^b	65.09 ^b	36.93 ^c	94.18 ^a	99.292	0.000
Nitrogen mineralization rate (mg kg ⁻¹ day ⁻¹)	0.31 ^a	0.08 ^b	-0.26 ^c	-0.24 ^c	95.401	0.000

*Results from the ANOVAs are included (F test and p value). Different letters in each line indicate significant differences ($p < 0.05$ by Duncan test) between forest stands

To select a representative MDS, the PCA method was used because of its MDS selection ability (Doran and Parkin, 1994). We performed standardized PCA of all data that showed statistically significant differences between different forest types via one-way analysis of variance (ANOVA) using the SPSS 19.0 statistical software package. The analytical hierarchy process (AHP) method was used to assign the data integration in an index (Lai *et al.*, 2002; Komac, 2006). An abstract view of such a hierarchy is shown in Figure 2.




Figure 2 A schematic diagram of the AHP process

In this research, Expert Choice software was used for determination of the best forest stand on basis of SQ indices using of AHP.

3 RESULTS

The soil characteristics having significant differences between the different forest types, thus, included for the PCA were: bulk density, silt, pH, EC, soil C/N, available nutrients (*i.e.*, P, K, Ca and Mg), earthworm biomass, microbial respiration, fine root biomass, nitrogen mineralization and litter C/N (Tables 1 and 2). The first and two component had Eigen value >1 (Table 2). The highly weighted variables under PC1 and PC2 were litter C/N ratio (-0.917), nitrogen mineralization (0.826), Ca (0.825), K (0.810), Mg (0.804), microbial respiration (0.785), EC (0.783), fine root biomass (0.759) and Silt (0.710), thus, were retained for the MDS (Table 2).

Table 2 PCA results of SQ indices having significant differences between the different forest stands

Principal components	PC1	PC2
Eigen value	6.843	1.371
Percent	48.879	9.791
Cumulative percent	48.879	58.669
	Eigen vectors- PC1	Eigen vectors- PC2
Bulk density (g cm ⁻³)	0.451	0.042
Silt (%)	-0.264	0.710
pH (1:2.5 H ₂ O)	0.664	0.078
EC (ds m ⁻¹)	0.783	0.111
Soil C/N	-0.623	-0.168
Available P (mg kg ⁻¹)	0.694	-0.145
Available K (mg kg ⁻¹)	0.810	0.141
Available Ca (mg kg ⁻¹)	0.825	0.106
Available Mg (mg kg ⁻¹)	0.804	-0.118
Earthworm biomass (mg m ⁻²)	0.635	0.023
Soil microbial respiration (mg CO ₂ -C g soil ⁻¹ day ⁻¹)	0.785	-0.107
Fine root biomass (g m ⁻²)	0.347	0.759
Litter C/N	-0.917	-0.060
Nitrogen mineralization rate (mg kg ⁻¹ day ⁻¹)	0.826	-0.400

Bold-italic factor loading correspond to the indicators included in the MDS

Different forest types were assessed, using of AHP approach, with respect to nine criteria of soil properties that were retained for the MDS (Figure 3). Inconsistency ratio values for every soil features in AHP are shown in Figures 3 and 4. According to our findings, the inconsistency ratios were less than 0.1 for whole of characters. Results are indicating that the maximum of local priority is belonging to *A. subcordata* on basis of EC, available nutrients (K, Ca and Mg), microbial respiration and N mineralization. Mixed natural forest had higher local priority with regarding to silt and fine root biomass characters, whereas the *T. distichum* had more local priority based on litter C/N (Figure 3). Determination of the criteria

role in assessment of different forest types and selection of the best forest stand as well as calculation of criteria weight were also carried out. For this purpose, the matrixes of paired comparisons were prepared and the criteria weights were calculated by arithmetic mean (Figure 4). Sensivity analysis is according to reported results also (Figure 5). The calculated overall priority showed that based on soil quality indices, the *A. subcordata* (0.370) forest type had higher ecological potential compared to the other stands. Whereas, *P. deltoids* (0.295), mixed natural forest (0.213), and *T. distichum* (0.122) had next priority, respectively (Table 3).




Figure 3 Local priority of different forest stands on basis of soil silt (a), EC (b), available K (c), available Ca (d), available Mg (e), microbial respiration (f), fine root biomass (g), litter C/N (h) and nitrogen mineralization (i)




Figure 4 Criteria priority based on arithmetic mean




Figure 5 Sensivity analysis based on performance alternative

Table 3 Overall priority of different forest stands on based on soil quality indices

Forest types	Overall priority	Class
<i>Alnus subcordata</i>	0.370	1
<i>Populus deltoids</i>	0.295	2
<i>Mixed natural forest</i>	0.213	3
<i>Taxadium distichum</i>	0.122	4

4 DISCUSSION

The study of SQ indices under each species was an approach to evaluate the performance of that species. Almost all of the nine indicators used with the MDS method can be found in

previously created MDS indicator methods (Doran and Parkin, 1994; Karlen and Stott, 1994; Larson and Pierce, 1994; Singer and Ewing, 2000; Ditzler and Tugel, 2002; Mastro *et al.*, 2008; Marzaioli *et al.*, 2010; Soleimany

Rahimabady et al., 2015). Among soil texture fractions, silt content was significantly higher under mixed natural forest in comparison to *T. distichum*, *P. deltoids* and *A. subcordata* plantation. This suggests a different evolution of the soil profile when covered by forest, especially in terms of erosion (Kooch et al., 2012 a, b).

The soil EC of the different species significantly follows the *A. subcordata* > *P. deltoids* > mixed natural forest > *T. distichum*. These differences may be caused by different foliage properties and the litter quality (Nsabimana et al., 2008; Haghdoost et al., 2011). Available K of soils of the *A. subcordata* was significantly higher than the soils of *P. deltoids*, mixed natural forest and *T. distichum*. In the present study, *A. subcordata* plantation with the highest K content can be attributed to rapid recycling of soil nutrients from trees (Chase and Singh, 2014). The lowest soil available K in *T. distichum* might be related to K retranslocation, nutrition and nutrient return (Rostamabadi et al., 2013). Hardwoods, including *A. subcordata*, typically contain a much greater amount of Ca^{2+} than to conifers, when growing on comparable sites (Ovington, 1956; Ovington, 1962). Soil Mg^{2+} had often been reported to be an element prone to leaching (Laskowski et al., 1995). The difference in the behavior of Mg may be due to the selective immobilization of nutrients by microbes (Rostamabadi et al., 2013). According to Hagen-Thorn et al. (2004), conifer species compared to hardwood stands, due to higher tendency to absorb base cations namely Mg makes reduced concentrations of these nutrients in the soil.

SMR was the lowest in *T. distichum* plantation as compared to other forest types. The highest SMR was found under *A. subcordata* plantation than can be related to the content of total N as an N-fixing species (Mo et al., 2004). Fine root biomass was significantly greater for the mixed natural forest than for the

plantations. Lee and Jose (2003) reported that hardwood forests had greater fine root biomass production compared to conifer forests. The big trees in the natural stand probably contributed to a comparatively higher fine root biomass (Tamooha et al., 2008). According to Nadelhoffer and Raich (1992), fine root production and aboveground production were linked with one another and were affected by similar factors. Height and diameter increments were measured to relate with the fine root production (Dipesh and Schuler, 2013). A similar influence of stand structure on root productivity has been reported in other studies (Le Goff and Ottorini, 2001; Hertel et al., 2009). The average D.B.H of trees was found to be a good predictor for fine root biomass and productivity (Drexhage and Colin, 2001; Le Goff and Ottorini, 2001; Hertel et al., 2009). Small diameter stems caused a significant reduction in fine root biomass (Joslin et al., 2000, Jones et al., 2003). Also different composition of tree species in the natural forest, compared with the monoculture plantations, may explain the higher fine root biomass in the mixed natural forest (Yang et al., 2003; Yang et al., 2004).

Valverde-Barrantes et al. (2014) find that most of the canopy tree species in their forest behave in largely the same way, with equal fine root proliferation in high resource patches. As a result of this species roots tend to aggregate in nutrient rich soils, resulting in a greater diversity of species within a given patch. They further find that fine root biomass and species diversity are greater in these soils (Jones, 2015). Dipesh and Schuler (2013) pointed that younger stands have less fine root production than the older stands. Fine roots could be easily affected by soil environmental factors (Eissenstat et al., 2000; Xu et al., 2013). The soil in *A. subcordata* and *P. deltoids* was more fertile and was in favor of the growth of fine root. The rich nutrients were in favor of fine root growth. The high fertility in the upper soil layer was a vital

factor affecting fine roots plantation (Wang *et al.*, 2014).

Nitrogen-fixing species plantations could increase soil nitrogen mineralization (Berg *et al.*, 2001; Rothe *et al.*, 2002). The difference in soil nitrogen mineralization was due to difference in availability of labile N substrates (McKinley *et al.*, 2008). High-quality litter decrease microbial immobilization of nitrogen, and result in enhanced N_{min} and plant available N (McKinley *et al.*, 2008). Van der Krift *et al.* (2001) found soil N mineralization of 5- and 15-year-old mixed poplar stands were greater than those of corresponding pure poplar stands. It has been reported that there was a negative relationship between the C/N and soil nitrogen mineralization in various ecosystems (Van der Krift *et al.*, 2001; Arslan *et al.*, 2010). We found a less nitrogen mineralization in forest types with higher C/N that is similar with Zeller *et al.* (2000) and Arslan *et al.* (2010) findings.

Litter C/N was the lowest in *A. subcordata* plantation as compared to other species and mixed natural forest. The narrow C/N was a precondition for the fast decomposition of organic matter (Ohta and Kumada, 1978). The C/N in the foliage of broadleaf species in the temperate zone is on average 25 (Vitousek *et al.*, 1988). C/N in organic litter was narrowed, because their dying led to the input of organic matter rich in nitrogen. Narrow C/N in the litter fall under *A. subcordata* monocultures caused the greatest part of carbon to be converted into CO₂ by oxidative processes as the end product of organic matter decomposition. Because of the decomposition of *A. subcordata* dead organic residues into end products, there was no intensive accumulation of organic C and N under *A. subcordata* plantations (Miletić *et al.*, 2012). Significantly higher C/N were found in the *T. distichum* stand these results underline the more recalcitrant nature of coniferous litter, probably due to the hard cuticle of needles, and could suggest a longer mean residence time of

this organic matter (Kooch *et al.*, 2012b). According to our findings, based on different SQ indices, the *A. subcordata* forest type had higher ecological potential compared to the other stands. Whereas, *P. deltoids*, mixed natural forest, and *T. distichum* had next priority, respectively.

5 CONCLUSION

Due to increasingly destruction of Hyrcanian Forests, plantation with native species was an appropriate method for rehabilitation and reconstruction of destroyed forest areas. To select a species, in addition to growth quantity and quality the effects of species in ecosystem, restoration should also be noted. Evaluation of trees in terms of soil moderator of different habitats conditions and classification of tree species based on achieved results were necessary. It appears using AHP gives a broad perspective in relation to the assessment of forest stands and can be considered an appropriate strategy. Using of AHP approach, as a conclusion, the N-fixing species, *A. subcordata*, is more efficient in improving soil quality in degraded land.

6 ACKNOWLEDGEMENT

Many thanks are due to Mr. Mohammad Beiranvand and Mr. Ali Khudadoost for their tireless assistance in field sampling. The authors are particularly grateful to Eng. Sadegh Boor for his expert technical assistance and to all the laboratory assistants at the faculty for analyzing the samples. This research was done by financial supports of Tarbiat Modares University, Iran, in the form of a Master science thesis of forestry.

7 REFERENCES

Alef, K. Estimating of soil respiration. In: "Methods in soil microbiology and biochemistry" (Alef K, Nannipieri P

- eds). Academic Press, New York, 1995; 464-470.
- Allison, L.E. Organic carbon In: "Methods of soil analysis" (Black CA ed). American Society of Agronomy, Part 2, Madison, WI, 1975; 1367-1378.
- Andrews, S.S., Karlen, D.L. and Cambardella, C.A. The soil management assessment framework. *Soil Sci. Soc. Am. J.*, 2004; 8: 1945-1962.
- Andrews, S.S., Karlen, D.L. and Mitchell, J.P.A. A comparison of soil quality indexing methods for vegetable production systems in Northern California. *Agric. Ecosyst. Environ.*, 2002; 90: 25-45.
- Aparicio, V. and Costa, J.L. Soil quality indicators under continuous cropping systems in the Argentinean Pampas. *Soil Tillage Res.*, 2007; 96: 155-165.
- Arshad, M.A. and Martin, S. Identifying critical limits for soil quality indicators in agroecosystems. *Agric. Ecosyst. Environ.*, 2002; 88: 153-160.
- Arslan, H., Guleryu, G. and Kirmızı, S. Nitrogen mineralization in the soil of indigenous oak and pine plantation forests in a Mediterranean environment. *Eur. J. Soil Biol.*, 2010; 46: 11-17.
- Asadiyan, M., Hojjati, S.M., Pourmajidian, M. R. and Fallah, A. Impact of land-use management on nitrogen transformation in a mountain forest ecosystem in the north of Iran. *J. For. Res.*, 2013; 24: 115-119.
- Ball, B.C., Batey, T. and Munkholm, L. J. Field assessment of soil structural quality - a development of the Peerkamp test. *Soil Use Manage.*, 2007; 23: 329-337.
- Berg, B., McClaugherty, C.A., De Santo, A.V. and Johnson, D.W. Humus buildup in boreal forests: effects of litter fall and its N concentration. *Can. J. Forest Res.*, 2001; 31: 988-998.
- Bouyoucos, G.J. Hydrometer method improved for making particle size analysis of soils. *J. Agron.*, 1962; 56: 464-465.
- Bower, C.A., Reitemeier, R.F. and Fireman, M. Exchangeable cation analysis of saline and alkali soils. *Soil Sci.*, 1952; 73: 251-261.
- Bremner, J.M. and Mulvaney, C.S. Nitrogen - total. In: "Methods of Soil Analysis" (page AL, Miller RH, Keeney RR eds). Second ed. American Society of Agronomy, Part 2, Madison, WI, 1982; 595-624.
- Caravaca, F., Figueroa, D., Barea, J.M., Azcón-Aguilar, C., Palenzuela, J. and Roldán, A. The role of relict vegetation in maintaining physical, chemical, and biological properties in an abandoned *stipa*-grass agro ecosystem. *Arid Land Res. Manage.*, 2003; 17: 103-111.
- Carrillo, Y., Ball, B. A., Strickland, M. S. and Bradford, M.A. Legacies of plant litter on carbon and nitrogen dynamics and the role of the soil community. *Pedobiologia*, 2012; 55: 185-192.
- Carter, M.R. Soil quality for sustainable land management. *J. Agron.*, 2002; 94: 38-47.
- Chase, P. and Singh, O.P. Soil nutrients and fertility in three traditional land use systems of Khonoma. *Res. Environ. J.*, 2014; 4: 181-189.
- Dipesh, K.C. and Schuler, J.L. Estimating fine-root production and mortality in the biomass plantations. *Comm. Soil Sci. Plant Anal.*, 2013; 44: 2514-2523.
- Ditzler, C.A. and Tugel, A.J. Soil quality field tools of USDANRCS soil quality institute. *Agron. J.*, 2002; 94: 33-38.

- Doran, J.W. and Parkin, T.B. Defining and assessing soil quality. In: Doran, J.W., Coleman, D.C., Bezdicek, D.F., Stewart, B.A. (Eds.), *Defining Soil Quality for a Sustainable Environment*. Soil Science Society of America, Madison, WI, USA, Special Publication, 1994; 35: 3-21.
- Drexhage, M. and Colin, F. Estimating root system biomass from breast-height diameters. *J. Forest.*, 2001; 74: 491-497.
- Dumanski, J. and Pieri, C. Land quality indicators: research plan. *Agric. Ecosyst. Environ.*, 2000; 81: 93-102.
- Eissenstat, D.M., Wells, C.E., Yanai, R.D. and Whitbeck, J.L. Building roots in a changing environment: implications for root longevity. *New Phytol.*, 2000; 147: 33-42.
- Govaerts, B., Sayre, K.D. and Deckers, J. A minimum data set for soil quality assessment of wheat and maize cropping in the highlands of Mexico. *Soil Tillage Res.*, 2006; 87: 163-174.
- Hagen-Thorn, A., Callesen, I., Armolaitis, K. and Nihlgard, B. The impact of six European tree species on the chemistry of mineral topsoil in forest plantations on former agricultural land. *For. Ecol. Manage.*, 2004; 195: 373-384.
- Haghdoust, N., Akbarinia, M., Hosseini, S. M. and Kooch, Y. Conversion of Hyrcanian degraded forests to plantations: Effects on soil C and N stocks. *Ann. Biol. Res.*, 2011; 2: 385-399.
- Herrick, J. E. Soil quality: an indicator of sustainable land management? *Appl. Soil Ecol.*, 2000; 15: 75-83.
- Hertel, D., Harteveld, M. A. and Leuschne, C. Conversion of a tropical forest into agroforestry alters the fine root-related carbon flux to the soil. *Soil Biol. Biochem.*, 2009; 41: 481-490.
- Homer, C. D. and Pratt, P. F. *Methods of analysis for soils, plants and waters*. University of California, Agricultural Sciences Publications, Berkeley, CA, 1961; 309P.
- Jones, A. Belowground fine root productivity, traits, and trees. *New Phytologist.*, 2015; 205: 461-462.
- Jones, R. H., Mitchell, R. J., Stevens, G. N. and Pecot, S. D. Controls of fine root dynamics across a gradient of gap sizes in a pine woodland. *Oecologia*, 2003; 134: 132-143.
- Jordan, D., Li, F., Ponder, F., Berry, E. C., Hubbard, V. C. and Kim, K.Y. The effects of forest practices on earthworm populations and soil microbial biomass in a hard wood forest in Missouri. *Appl. Soil Ecol.*, 1999; 13: 31-38.
- Joslin, J. D., Wolfe, M. H. and Hanson, P. J. Effects of altered water regimes on forest root systems. *New Phytologist.*, 2000; 147: 117-129.
- Karlen, D. L. and Scott, D. E. A framework for evaluating physical and chemical indicators of soil quality. In: Doran, J.W., Coleman, D.C., Bezdicek, D.F., Stewart, B.A. (Eds.), *Defining Soil Quality for a Sustainable Environment*. ASA and SSSA, Madison, WI, USA, 1994; 53-72.
- Karlen, D. L., Andrews, S. S., Doran, J. W. and Wienhold, B. J. Soil quality: humankind's foundation for survival. *J. Soil Water Conserv.*, 2003; 58: 171-179.
- Komac, M. A landslide susceptibility model using the analytical hierarchy process method and multivariate statistics in per

- alpine Slovenia. *Geomorphology*, 2006; 74: 17-28.
- Kooch, Y., Hosseini, S. M., Mohammadi, J. and Hojjati, S. M. Determination of the Best Canopy Gap Area on the Basis of Soil Characteristics Using of Analytical Hierarchy Process (AHP). *Folia For. Polon.*, 2012a; 54: 15-24.
- Kooch, Y., Hosseini, S. M., Zaccone, C., Jalilvand, H. and Hojjati, S. M. Soil organic carbon sequestration as affected by afforestation: the Darab Kola forest (North of Iran) case study. *J. Environ. Monit.*, 2012b; 14: 2438-2446.
- Kooch, Y., Theodose, T. A. and Samonil, P. Role of deforestation on spatial variability of soil nutrients in a Hyrcanian forest. *Ecopersia*, 2014; 2: 779-803.
- Lafleur, B., Labrecque, M., Arnold, A. A. and Bélanger, N. Organic carbon accumulation in topsoil following afforestation with Willow: emphasis on leaf litter decomposition and soil organic matter quality. *Forests*, 2015; 6: 769-793.
- Lai, V. S., Wong, B. K., Cheung, W. Group decision making in a multiple criteria environment: a case using the AHP in software selection. *Eur. J. Oper. Res.*, 2002; 137: 134-144.
- Larson, W. E. and Pierce, F.J. The dynamics of soil quality as a measure of sustainable management. *Defining Soil Quality for a Sustainable Environment*. Soil Science Society of America, Madison, Wisconsin, 1994; 37-52. USA.
- Laskowski, R., Nikiski, M. and Maryanska, M. The dynamics of chemical elements in forest litter. *Ecology*, 1995; 76: 1393-1406.
- Le Goff, N. and Ottorini, J. M. Root biomass and biomass increment in a beech (*Fagus sylvatica*) stand in North East France. *Ann. For. Sci.*, 2001; 58: 1-13.
- Lee, K. H. and Jose, S. Soil respiration, fine root production, and microbial biomass in cottonwood and loblolly pine plantations along a nitrogen fertilization gradient. *For. Ecol. Manage.*, 2003; 185: 263- 273.
- Li, G., Chen, J., Sun, Z. and Tan, M. Establishing a minimum dataset for soil quality assessment based on soil properties and land-use changes. *Acta Ecol. Sin.*, 2007; 27: 2715-2724.
- Lima, A. C., Brussaard, L., Totola, M. R., Hoogmoed, W. B. and de Goede, R. G. A functional evaluation of three indicator sets for assessing soil quality. *Appl. Soil Ecol.*, 2013; 64: 194-200.
- Marzaioli, R., D'Ascoli, R., De Pascale, R. A. and Rutigliano, F. A. Soil quality in a Mediterranean area of Southern Italy as related to different land use types. *Appl. Soil Ecol.*, 2010; 44: 205-212.
- Masto, R., Chhonkar, P., Singh, D. and Patra, A. Alternative soil quality indices for evaluating the effect of intensive cropping, fertilization and managing for 31 years in the semi-arid soils of India. *Environ. Monit. Assess.*, 2008; 136: 419-435.
- Mc Kinley, D.C., Rice, C.W. and Blair, J.M. Conversion of grassland to coniferous woodland has limited effects on soil nitrogen cycle processes. *Soil Biol. Biochem.*, 2008; 40: 2627-2633.
- Miletic, Z., Knezevic, M., Stajic, S., Kosanin, O. and Dordevic, I. Effect of European black Alder monocultures on the characteristics of reclaimed mine soil. *Inter. J. Environ. Res.*, 2012; 6: 703-710.

- Mo, J.M., Xue, J.H., and Fang, Y.T. Litter decomposition and its responses to simulated N deposition for the major plants of Dinghushan forests in subtropical China. *Acta Ecol. Sin.*, 2004; 24: 1513-1420.
- Nadelhoffer, K.J. and Raich, J.W. Fine root production estimates and belowground carbon allocation in forest ecosystems. *Ecology*, 1992; 73: 1139-1147.
- Neatrou, M.A., Jones, R.H. and Golladay, S. W. Correlations between soil nutrients availability and fine - root biomass at two spatial scales in forested wetlands with contrasting hydrological regimes. *Can. J. For. Res.*, 2005; 35: 2934-2941.
- Nsabimana, D., Klemedtson, L., Kaplin, B. A. and Wallin, G. Soil carbon and nutrient accumulation under forest plantations in southern Rwanda. *African J. Environ. Sci. Tech.*, 2008; 2: 142-149.
- Ohta, S. and Kumada, K. Studies on the humus forms of forest soils. VI. Mineralization of nitrogen in brown forest soils. *Soil Sci. Plant Nut.*, 1978; 24: 41-54.
- Ovington, J.D. Quantitative ecology and the woodland ecosystem concept. In J. B. Cragg (ed.), *Advances in ecological research*. Academic Press, London and New York: Elsevier. 1962; 103-192.
- Ovington, J.D. Studies of the development of woodland conditions under different trees. V. The mineral composition of the ground flora. *J. Ecol.*, 1956; 42: 597-604.
- Plaster, E.J. *Soil Science and Management*. Delmar Publishers Inc., Albany, NY., 1985; 124 P.
- Poorzady, M. and Bakhtiari, F. Spatial and temporal changes of Hyrcanian forest in Iran. *For.*, 2009; 2: 198-206.
- Qi, Y., Darilek, J.L., Huang, B., Zhao, Y., Sun, W. and Gu, Z. Evaluating soil quality indices in an agricultural region of Jiangsu Province, China. *Geoderma*, 2009; 149: 325-334.
- Rezaei, S.A., Gilkes, R.J. and Andrews, S.S. A minimum data set for assessing soil quality in rangelands. *Geoderma*, 2006; 136: 229-234.
- Rostamabadi, A., Tabari, M. and Sayad, E. Influence of *Alnus subcordata*, *Populus deltoides* and *Taxodium distichum* on poor drainage soil, northern Iran. *Ecopersia*, 2013; 1: 207-218.
- Rothe, A., Cromack, J.K., Resh, S.C., Makeneci, E. and Son, Y. Soil carbon and nitrogen changes under Douglas-fir with and without red alder. *Soil Sci. Soc. Am. J.*, 2002; 66: 1988-1995.
- Sagheb-Talebi, Kh., Sajedi, T. and Pourhashemi, M. *Forests of Iran: A treasure from the past, a hope for the future*. Springer Publications, 2014; 149 p.
- Shepherd, T.G. *Visual soil assessment, field guide for pastoral grazing and cropping on flat to rolling country*. Horizons Regional Council. Palmerston North, 2009; 1: 119 P.
- Singer, M.J. and Ewing, S. Soil quality. In: Sumner, M.E. (Ed.), *Handbook of Soil science*. CRC Press, Boca Raton, FL, USA, 2000; G-271-G-298.
- Soleimany Rahimabady, M., Akbarinia, M. and Kooch, Y. The effect of land covers on soil quality properties in the Hyrcanian regions of Iran. *J. Bios. Biotech.*, 2015; 4: 73-79.
- SPSS Inc. Released. *SPSS for Windows, Version 19.0*. Armonk, NY, SPSS Inc. 2010.

- Tamooha, F., Huxhamd, M., Karac, M., Mencuccinie, M., Kairoc, J. G. and Kirui, B. Below-ground root yield and distribution in natural and replanted mangrove forests at Gazi bay. *For. Ecol. Manage.*, 2008; 256: 1290-1297.
- Valverde-Barrantes, O.J., Smemo, K.A., Feinstein, L.M. and Kershner, M.W. Aggregated and complementary: symmetric proliferation, over yielding, and mass effects explain fine root biomass in soil patches in a diverse temperate deciduous forest landscape. *New Phytologist*, 2014; 205: 731-742.
- Van der Krift, T.A.J., Gioacchini, P., Kuikman, P. J. and Berendse, F. Effects of high and low fertility plant species on dead root decomposition and nitrogen mineralization. *Soil Biol. Biochem.*, 2001; 33: 2115-2124.
- Vitousek, P.M., Fahey, T., Johnson, D.W. and Swift, M.J. Element interactions in forest ecosystems: succession, allometry, and input-output budgets. *Biogeochemistry*, 1988; 5: 7-34.
- Wang, X., MA, L., JIA, Zh, and Jia, L. Root inclusion net method: novel approach to determine fine root production and turnover in *Larix principis-rupprechtii* Mayr plantation in North China. *Turk. J. Agri. For.*, 2014; 38: 388-398.
- Xu, W., Liu, J., Liu, X., Kun Li, K., Zhang, D. and Yan, J. Fine root production, turnover, and decomposition in a fast-growth *Eucalyptus urophylla* plantation in southern China. *J. Soils Sed.*, 2013; 13: 1150-1160.
- Yang Y.S., Guo, J.F., Chen, G.S., He, Z. M. and Xie, J. S. Effects of slash burning nutrient removal and soil fertility in Chinese fir and evergreen broadleaved forests of mid-subtropical China. *Pedosphere*, 2003; 13: 87-96.
- Yang, Y.S., Chen, G.S., Lin, P., Xie, J.S. and Guo, J. F. Fine root distribution, seasonal pattern and production in four plantations compared with a natural forest in Subtropical China. *Ann. For. Sci.*, 2004; 61: 617-627.
- Zeller, V., Bahn, M., Aichner, M., and Tappeiner, U. Impact of land-use change on nitrogen mineralization in the Southern Alps. *Biol. Fert. Soils*, 2000; 31: 441-448.
- Zhang, B., Zhang, Y., Chen, D., White, R. E. and Li, Y. A quantitative evaluation system of soil productivity for intensive agriculture in China. *Geoderma*, 2004; 123: 319-331.
- Zhang, C., Xue, S., Liu, G.B. and Song, Z.L. A comparison of soil qualities of different vegetation types in the Loess Plateau, China. *Plant Soil*, 2011; 347: 163-178.

شاخص‌های کیفیت خاک در توده‌های جنگلی خالص و آمیخته منطقه جنوبی دریای خزر

یحیی کوچ^{۱*}، فاطمه روستایی^۲ و سید محسن حسینی^۳

۱- استادیار، دانشکده منابع طبیعی، دانشگاه تربیت مدرس، ایران

۲- دانشجوی کارشناسی ارشد، دانشکده منابع طبیعی، دانشگاه تربیت مدرس، ایران

۳- استاد گروه جنگلداری، دانشکده منابع طبیعی، دانشگاه تربیت مدرس، ایران

تاریخ دریافت: ۹ فروردین ۱۳۹۴ / تاریخ پذیرش: ۱۲ مرداد ۱۳۹۴ / تاریخ چاپ: ۹ آبان ۱۳۹۴

چکیده پژوهش حاضر با هدف ارزیابی گونه‌های خالص کاشته شده (شامل توسکا ییلاقی، صنوبر دلتوئیدس، دارتالاب) و توده طبیعی آمیخته (با غالبیت گونه‌های بلوط- ممرز- انجیلی) بر مبنای برخی شاخص‌های کیفیت خاک در استان مازندران، شمال ایران، صورت پذیرفت. در هر یک از توده‌های مورد بررسی، تعداد ۱۶ نمونه خاک (۱۰-۰ سانتی‌متری) برداشت و مشخصه‌های چگالی ظاهری، بافت، محتوی رطوبت، pH، هدایت الکتریکی، کربن آلی، نیتروژن کل، عناصر غذایی قابل جذب، زیتوده کرم‌های خاکی، تنفس میکروبی، زیتوده ریزریشه‌ها، کربن و نیتروژن لاشبرگ در محیط آزمایشگاه اندازه‌گیری شد. از بین متغیرهای مورد بررسی، نه معیار (سیلت، هدایت الکتریکی، پتاسیم، کلسیم، منیزیم، تنفس میکروبی، زیتوده ریزریشه، معدنی شدن نیتروژن، نسبت کربن به نیتروژن لاشبرگ) بر مبنای تحلیل مؤلفه‌های اصلی به عنوان حداقل مجموعه داده‌ها انتخاب شدند. به منظور ادغام مجموعه داده‌ها در یک شاخص، تحلیل فرآیند سلسله مراتبی (AHP) به کار گرفته شد. وزن نهایی محاسبه شده بر مبنای نه معیار نشان داد که توده جنگلی توسکا دارای توان اکولوژیکی (۰/۳۷۰) بالاتری نسبت به توده‌های دیگر می‌باشد و توده‌های صنوبر طبیعی آمیخته و دارتالاب به ترتیب با توان اکولوژیکی ۰/۲۹۵، ۰/۲۱۳ و ۰/۱۲۲ در مراحل بعدی قرار می‌گیرند. به‌عنوان نتیجه‌گیری، می‌توان اظهار نمود که گونه تثبیت‌کننده ازت، توسکا ییلاقی، دارای بازدهی بیش‌تری برای بهبود کیفیت خاک در مناطق جنگلی تخریب‌یافته داشته است.

کلمات کلیدی: جنگل‌های هیرکانی، گونه‌های پهن‌برگ، گونه‌های سوزنی‌برگ، مرکز بذر جنگلی خزر، مشخصه‌های خاک