



## The Interaction Effects of Allelopathy and Salinity Stresses on Physiological Traits and Nutrient Uptake in *Poa bulbosa* L. Treated with Arbuscular Mycorrhizal Fungi

### ARTICLE INFO

#### Article Type

Original Research

#### Authors

Karami F.<sup>1</sup> MSc,  
Akhzari D.\*<sup>1</sup> PhD

#### How to cite this article

Karami F, Akhzari D. The Interaction Effects of Allelopathy and Salinity Stresses on Physiological Traits and Nutrient Uptake in *Poa bulbosa* L. Treated with Arbuscular Mycorrhizal Fungi. ECOPERSIA. 2020;8(3):139-146.

<sup>1</sup>Nature Engineering Department, Natural Resources & Environmental Science Faculty, Malayer University, Malayer, Iran

#### \*Correspondence

Address: Natural Resources & Environmental Science Faculty, Malayer University, Malayer, Iran. Postal Code: 6571995863  
Phone: +98 (81) 32355330  
Fax: +98 (81) 33339840  
d\_akhzari@yahoo.com

#### Article History

Received: December 29, 2019

Accepted: February 09, 2020

ePublished: September 20, 2020

### ABSTRACT

**Aims** Allelopathy and salinity stresses are known as major abiotic factors which limits the growth and production of plants. Environmental stresses such as salinity and allelopathy can affect the growth and establishment of rangeland species; however the interaction of such environmental stress and inter-plant competition (i.e., allelopathic effects) associated with arbuscular mycorrhiza rarely understood. Therefore, this study aimed to investigate the effect of arbuscular mycorrhiza on physiological traits and nutrient uptake of *Poa bulbosa* L. under the influence of allelopathy and salinity stress.

**Materials & Methods** A factorial experiment was designed with a completely randomized design with three replications. Factors include allelopathy extracted by *Artemisia persica* Boiss root (treat vs. control), salinity (control, 1, 4, and 8mmol/kg NaCl) and mycorrhizal fungi (mixed and unmixed). The experiment was set up in Malayer University greenhouse in a completely randomized design with factorial layout and 3 replications.

**Findings** The stem and root of the studied plant species were higher than those found in non-inoculated treatments at low salinity and allelopathy treatments ( $p < 0.05$ ). Also, in severe stress conditions, mycorrhizal dependency was significantly higher (50.87). Proline content does not show variation under all studied treatments ( $p < 0.05$ ). Chlorophyll content decreased significantly with increasing salinity and allelopathy, but in the presence of mycorrhizal fungi, the rate of reduction was significantly lower and mycorrhizal fungi increased chlorophyll well. Plant nitrogen and phosphorus contents also coincided with the mycorrhizal fungi, showing the highest yield (3.7% and 2.6g kg<sup>-1</sup> for total nitrogen and phosphorus respectively).

**Conclusion** The growth, establishment and physiological properties of the *Poa bulbosa* L. are improved under the influence of mycorrhizal application. Thus, the use of mycorrhizal fungi can be a useful biological method to reduce the destructive effects of environmental stresses such as salinity and allelopathy.

**Keywords** Arbuscular mycorrhizal fungi; Allelopathy; Salinity stress; *Poa bulbosa* L.

### CITATION LINKS

[1] Accounting for the world's ... [2] Disturbances impact on longevity of grass seeds, semi-... [3] Allelopathic interactions between ... [4] Influence of coumarin on some physiological and biochemical ... [5] Soil salinity delays germination and limits growth of hyphae from propagules ... [6] Influence of rootstocks on the resistance of cashew ... [7] Phylogenetic distribution and evolution of ... [8] Dual colonization of Eucalyptus urophylla ST Blake by arbuscular and ectomycorrhizal ... [9] The effects of soil phosphorus and zinc availability on plant responses to mycorrhizal fungi: a ... [10] Mineral acquisition by arbuscular ... [11] Influence of arbuscular mycorrhizal mycelial exudates on soil bacterial ... [12] Some Root Traits of Barley (*Hordeum vulgare* L.) as Affected by Mycorrhizal ... [13] Evaluation of growth potential in spinach inoculated with *Piriformospora indica*, Mycorrhiza and dark ... [14] Improvement of salt tolerance in rice plants by arbuscular ... [15] Impact of two arbuscular mycorrhizal fungi on *Arundo* ... [16] Alleviation of salt stress and changes in glycyrrhizin accumulation by arbuscular ... [17] Alfalfa (*Medicago sativa* L.) clones tolerant to salt ... [18] Rapid determination of free proline ... [19] Effects of arbuscular mycorrhizal fungi on seedling growth and physiological ... [20] Studying the effect of mycorrhiza on root morphology and rhizosphere's pH ... [21] Studying the effects of salinity stress on the growth of various ... [22] Studying the effects of mycorrhiza and vermicompost fertilizers on the ... [23] The interaction of *Artemisia persica* allelopathy, drought and arbuscular mycorrhizal ... [24] Effect of extract of fast growing species *Trifolium alexandrinum* L ...