



Effect of Different Nutritional Treatments on the Growth of *Datura Stramonium* L.

ARTICLE INFO

Article Type Original Research

Authors

Esmaeil Sheidai-Karkaj, Ph.D.^{1*}
Esfandiar Jahantab Ph.D.²
Samira Sasanifar, Ph.D.³
Morteza Mofidi-Chelan, Ph.D.¹
Sara Babazadeh, M.Sc.⁴

How to cite this article

Sheidai-Karkaj E, Jahantab E, Sasanifar S, Mofidi-Chelan M. Babazadeh S. Effect of Different Nutritional Treatments on the Growth of *Datura Stramonium* L. ECOPERSIA 2025;13(1): 37-48.



10.22034/ECOPERSIA.13.1.37

¹ Associate Professor, Department of Range and Watershed Management, Faculty of Natural Resources, Urmia University, Urmia, Iran.

² Associate Professor, Department of Range and Watershed Management (Nature Engineering), Faculty of Agriculture, Fasa University, Fasa, Iran.

³ Former Ph.D. Student of Forestry, Department of Forestry, Faculty of Natural Resources, Urmia University, Urmia, Iran

⁴ M.Sc. Student of Rangeland Sciences and Engineering- Rangeland Management, Faculty of Natural Resources, Department of Range and Watershed Management, Urmia University, Urmia, Iran.

* Correspondence

Address: West Azerbaijan Province, Urmia, 11 km road of Serow, Nazlu Pardis, Faculty of Natural Resources
Tel: +00989144016823
Email: e.sheidai@urmia.ac.ir

Article History

Received: November 26, 2024
Accepted: February 1, 2025
Published: February 28, 2025

ABSTRACT

Aims: *Datura stramonium* L. is a medicinal plant known for its alkaloid compounds. Limited research has explored the impact of fertilizers like solopotas and fulvic acid on its growth and structural traits. This study aimed to evaluate the effects of these fertilizers on the growth rate and biomass of *D. stramonium*.

Materials & Methods: Solopotas and fulvic acid were applied as foliar sprays at concentrations of 2%, 4%, 6%, and 8%, with five applications at 5-day intervals. Growth parameters such as plant height, leaf length, stem length, root length, leaf number, flower and fruit counts, and shoot weight were measured at maturity and compared with control plants treated with distilled water.

Findings: The fertilizer treatments significantly affected leaf traits, flower and fruit numbers, plant height, and shoot weight ($p < 0.05$) but not root length. The 2% fulvic acid treatment resulted in the tallest plants (45.75 cm), compared to 31.25 cm in the control. The 4% fulvic acid treatment had the most leaves (39.5), while the control had the lowest (19). The 2% solopotas treatment produced the most extended leaves (14.15 cm), while the 8% solopotas had the shortest (8.17 cm). The 2% fulvic acid also resulted in the heaviest shoots (13.5 g), compared to 3.5 g in the control.

Conclusion: The application of 2% fulvic acid and 2% solopotas significantly improved the growth and biomass of *D. stramonium*, particularly in plant height, leaf number, and leaf length. These findings suggest that these fertilizers can enhance the commercial potential of *D. stramonium*.

Keywords: Solopotas and Fulvic Acid Fertilizer; Biomass; Medicinal Plant; Structural Traits.

CITATION LINKS

[1] Foroghi A. Review of ... [2] Beyk-Khormizi A., Sarafranz-Ardakani M.R., Hosseini Sarghein S., Mosh ... [3] Da Cunha Honorato A., de Assis R.M.A., Maciel J.F.A., Nohara G.A., d ... [4] Hao D., Luan Y., Wang Y., Xiao P. Unveiling Nitrogen Fertilizer in M ... [5] Ibrahim M.H., Jaafar H.Z., Karimi E., Ghasemzadeh A. Impact of organ ... [6] Oseni O., Olarinoye C., Amoo I. Studies on chemical compositions and ... [7] Niakan M., Rezapour Mahjoob S., Ghorbanli M. Effect of exogenous put ... [8] Al-Snafi A.E. Medical importance of *Datura fastuosa* (syn: *Datura met* ... [9] Singh L.R., Singh O.M. *Datura stramonium*: An overview of its phytoch ... [10] Ashtiania F., Sefidkonb F. ... [11] Heidarzadeh A., Mohammad Modarres-Sanavy S.A., Mokhtassi-Bidgoli A. ... [12] Mohammadpour Vashvaei R., Ghanbari A., Asgharipour M.R., Ramroudi M. ... [13] Izadi Z., Biabani A., Sabouri H., Bahreyninejad B. Effects of nitrog ... [14] Afsharypuor S., Mostajeran A., Shانه-Saz M. Effect of urea fertiliz ... [15] Abbaspour R., Imam Y., Hosseini S.Z. Investigating the effect of org ... [16] Hadinezhad F., Salimi A., Chavoushi M. Evaluation of the effect of n ... [17] Mardi M., Abbasifar A., ValizadehKaji B. Comparison of the effect of ... [18] Ehsani S., Niknahad Gharmakher H., Motamedi J., Akbarlou M., Sheidai ... [19] Najafzadeh Nobar Z., Safari Sinegani A.A. Effect of soil contaminati ... [20] Abbasi Khalaki M., Ghorbani A., ... [21] Gorji Z., Mehrparvar M., Mozaffari H. Organic vs. Inorganic Fertiliz ... [22] Seilsepour M., ... [23] Zhao Y., Wei Y., Zhang Y., Wen X., Xi B., Zhao X., Wei Z. Roles of c ... [24] Patnaik P. Handbook of inorganic chemicals: McGraw-Hill New York; 20 ... [25] Piccolo A. The Supramolecular structure of ... [26] Sasanifar S., Alijanpour A., Shafiei Banj A., Eshaghi Rad J.E., Mola ... [27] Hair Junior J.F., Black W.C., ... [28] Sheidai-Karkaj E., Sepehry A., ... [29] Arvin P. Study of ... [30] Makkizadeh M., Chaichi M., Nasrollahzadeh S., Khavazi K. Effect of d ... [31] Mollashahi M., Ganjali H., Fanaei H. Effect of different levels of n ... [32] Hossain M.D., Musa M.H., Talib J., Jol H. Effects of nitrogen, phosph ... [33] Fageria N.K. The use of nutrients in ... [34] Tehlan S., Thakral K., ... [35] Thalooth A., Tawfik M., Mohamed H.M. A comparative study on the effe ... [36] Alizad L., Mostafavi Rad ... [37] Abd Allah M.M., El-Bassiouny H.M., Elewa T.A.E., El-Sebai T.N. Effec ... [38] Mousavi Fazl S.H., ... [39] Habib H.A., Shafeek M., Zaki M., El-Shal Z. Response of potato plant ... [40] Rahimi M.M., ... [41] Lin S., Sattelmacher B., Kutzmutz E., Mühling K.H., Dittert K. Infl ...

Introduction

In addition to their aesthetic value and productivity, many plants have medicinal properties. Identifying these plant species and understanding the impact of environmental and nutritional factors on the concentration of active compounds, their chemical composition, and plant growth is both important and beneficial. Nutritional substances sometimes effectively strengthen these plants' medicinal aspects [1]. The use of medicinal plants as a source of healing in local communities worldwide has a history of thousands of years. Today, the role of medicinal plants as a primary healthcare method for most communities is not hidden. Medicinal plants have vast potential in treating various diseases due to the presence of important therapeutic phytochemicals. Therefore, pharmaceutical sources are very important, so a significant share of all synthetic drugs comes from them. Iran has a special place in the world regarding geographical location, climate, and weather, and it is the habitat of many plant species, including medicinal plants. Climatic diversity in Iran has caused the existence of few medicinal plants in the world that do not have the same or similar species in Iran. Various researches have shown that it is possible to achieve high production due to suitable conditions for growing medicinal plant species [2, 3, 4, 5]. On the other hand, due to the growing economic importance of medicinal plants and also because most of the medicinal plants are native to developing countries, it is possible to provide part of the significant health and treatment needs of the society with proper use.

Datura stramonium, an annual plant known by common names such as thornapple, jimsonweed, and devil's trumpet, is a toxic flowering species in the Daturae tribe of the Solanaceae family. Its likely origin is in Central America and has been

introduced in many world regions [6]. This plant is abundantly observed in the coastal areas of northern Iran from Astara to the east of Mazandaran [7]. *D. stramonium* is one of the most important medicinal plants [8, 9]. This plant has antibacterial, insecticidal, repellent, antioxidant, anti-inflammatory, analgesic and antipyretic, antispasmodic, anticancer, antifertility, sedative, antiasthmatic, invigorating and protective against organophosphate toxins [8, 9]. Also, this plant is used to treat diabetes, asthma, herpes, diarrhea, menstrual cramps, painful skin ulcers, stomach inflammation, headache, and muscle cramps [8, 9]. The important phytochemical compounds in *D. stramonium* are alkaloids, and a group of alkaloids called tropane alkaloids are mainly found in plants of the Solanaceae family, especially in the *Datura* genus [10].

The growth and performance of plants are influenced by genetic and environmental factors, which are essential for expressing a plant's genetic traits. Plant growth will be disrupted or halted if either of these factors is lacking or deficient. Properly using agricultural methods and factors in medicinal plant production is essential to increase the number of effective substances [11]. In this regard, it seems necessary to investigate detailed scientific research to determine the effect of different fertilizers on the growth rate and performance of medicinal plants. In the following section, various research studies that have been done on different plants will be reviewed. In a study, Mohammadpour Vashvaei et al. (2021) evaluated the physiological, biochemical, and agronomic characteristics of *D. stramonium* under the residual effect of conventional and low-input cultivation systems. The results showed that the highest amount of plant height, number of branches per plant, number of leaves per plant, and dry weight of leaves belonged to the combined

treatment of vermicompost and nanobiomic fertilizers ^[12]. In research, the researchers investigated the effect of different levels of nitrogen and density on the growth and performance of the *D. stramonium* plant. The results showed that the application of 180 kg. ha⁻¹ of urea and the density of 6 plants per unit area can be essential in increasing the seed yield of the medicinal plant *D. stramonium* ^[13]. Investigating the effect of different levels of urea fertilizer on *D. stramonium* showed that the use of 240 kg. ha⁻¹ of urea increases the biomass yield and grain yield [14]. In a study using 150 and 300 kg. ha⁻¹ of nitrogen fertilizer on the *D. stramonium* plant, the results showed that the treatment of 300 kg. ha⁻¹ increased the growth and the percentage of seed germination in the *D. stramonium* plant ^[15]. Hadinezhad et al. (2021) stated that using biological and combined fertilizers (biological and chemical) and providing the necessary materials to increase the growth of *Nigella sativa* L. is effective. It will lead to better efficiency ^[16]. Mardi et al. (2022) reported that the use of fertilizers such as supernitroplus has increased plant height, functional traits, and the number of leaves compared to control (no fertilizer use) in *Allium sativum* L. ^[17]. Ehsani et al. (2021) surveyed the effect of lignite and wheat straw biochar on vegetation growth for *Astragalus podolobus* Boiss. & Hohen found that with the addition of lignite to the soil, the maximum height, crown diameter, canopy cover, and volume of biomass were recorded for a 3.75% application rate after the fourth month and were 25.03cm, 23.52cm, 528.65cm², and 15581cm³, respectively ^[18]. Najafzadeh Nobar and Safari Sinegani (2020) reported that the antioxidant activity of green bean biomass was improved in mine soil in comparison with rangeland and cropland soils (37.76% and 18.43%, respectively) ^[19]. Abbasi Khalaki et al. (2021)

concluded that the maximum plant viability (80, 82.22%), height (59, 68.33cm), leaf area index (67.79, 84.93cm²), basal diameter (13.33, 16.66cm), canopy cover (993.33, 1242.66cm²), relative water content (70.46, 88.32%), photosynthesis rate (27.29, 28.49μmolco₂.cm⁻².s⁻¹), chlorophyll (0.87, 0.72mg.g⁻¹), number of inflorescence (57.33, 56Nm⁻²) and 1000-grain weight (33.30, 3.89g.m⁻²) were in sainfoin (EM2%) and alfalfa (PSN1000mg lit⁻¹), respectively ^[20]. Gorji et al. (2023), in comparison of the impact of organic and inorganic fertilizers on phenol content and growth traits in cultivated and wild safflower, found that the leaf area, number of leaves, dry and fresh biomass weights significantly increased in the case of adding vermicompost to the soil ^[21]. Seilsepour and Moharami (2024) surveyed the impact of nitrogen adding and municipal solid waste compost on wheat yield in soil with high caco₃. The experiment's results demonstrated that the effect of MSW compost with N-fertilizer and their interaction on grain yield and biological function were significant ^[22].

Considering the high value and importance of the medicinal plant *D. stramonium* and on the other hand, since there has been no report on the effect of solopotas and fulvic fertilizers on the structural characteristics and biomass of the *D. stramonium* plant, it seems necessary to investigate the impact of these treatments on the medicinal plant *D. stramonium*. Therefore, considering the importance of *D. stramonium* as a medicinal plant and its widespread use in the pharmaceutical industry, the present research was conducted to survey the effects of different treatments on the structural traits and biomass of *D. stramonium*. The most important purpose and context of this research is to support the necessities for plant domestication. Since most medicinal plants, such as *Datura stramonium* L., are

suitable for producing drug products, several efforts are needed to domesticate them. On the other hand, the vast area of dry farming lands is abandoned, and then it is required to be restored. This species is one of the candidate species for cropping in this area.

Materials & Methods

In the present study, the seeds of the *D. stramonium* plant were collected from the region of Nazlu in Urmia, West Azerbaijan Province, and the dried seeds of this plant were removed and used. Because of controlling the effects of environmental factors on germination and growth, the pots were filled by the soil of the *D. stramonium* species habitat in the Nazlu region. The soil texture was loamy-clay. Its pH was about 7.1, and EC was 0.23 $\text{mhos}\cdot\text{cm}^{-1}$. The research started on January 1, 2021, and the seeds were first placed in the cultivation trays and watered every other day. After the plants reached 10-12 cm in height, they were transferred to larger pots, and the same watering routine was resumed. After the primary growth of the plants, the treatment was started. In this way, two types of fertilizer, solopotas, and fulvic acid, were used. Fulvic acid is used as a humic substance resulting from the decomposition of the remains of living organisms, without harmful environmental effects, in the production of agricultural and garden crops. Fulvic acid is a powerful organic electrolyte that regulates and energizes cells [23]. By entering the plant tissue, fulvic acid brings macro elements with it from the surface part of the plant to the plant tissue [24]. Also, potassium sulfate fertilizer or solopotas with 51% potassium and 17.5% sulfur is the best source of potassium, and due to its sulfur base, it can be easily used in salty soils and has the least leaching. This substance is obtained from the combination of potassium oxide K_2O 51% and sulfur 46%

SO_3 and is very widely used; this fertilizer is one of the most harmless chemical and inorganic fertilizers. Potassium plays a role in transporting carbohydrates and other micronutrients in the stem and leaves. On the other hand, strengthening the plant and stem makes it more resistant to cold, heat, pests, and diseases. Solopotas is entirely soluble in water and allows foliar spraying without creating deposits in the nozzles. Solopotas fertilizer increases the nutritional value of products by positively affecting the production of starch, sugar, and vitamins [25]. The treatment was done by spraying solutions with different doses of fertilizers on the plant's leaves, buds, and flowers without falling on the soil's surface. A total of 27 pots of plants were available, and three replications were considered for each percentage of fertilizer. Thus, three pots were used as control plants: three for 2%, 4%, 6%, and 6%, and solopotas for 8%. The same procedure was repeated for fulvic acid. The pots were placed in the greenhouse of Urmia University with the same temperature and humidity conditions, and the treatment was done five times with an interval of 5 or 6 days (Figure 1). Irrigation was carried out every other day to prevent its level from decreasing to less than the field capacity (FC) during the experiment. After 35 days, the height and size of the plants were measured before separating the organs in question. This way, the length from the soil surface to the highest part of the plant was recorded as the height. The number of the smallest to the largest leaves was counted and recorded as the number of leaves. In the same way, the number of fruits, flowers, buds, and the length and width of two leaves from each pot were also noted, and the crown of the plant, the length of its large and small crowns were also reported, and after removing the plants from the pot, the length of the root was also recorded.



Figure 1) *D. stramonium* plant in the research greenhouse.

The parameters considered during the measurement are flower, leaf, stem, root weight in grams, small crown length, considerable crown length, leaf width, leaf length, number of flowers and fruits, number of leaves, root length, and plant height. The measurements were accurately taken using a tape measure.

Analysis Method

In the current study, nine treatments, including solopotas solution spray at four concentration levels of 2, 4, 6, and 8% and fulvic acid solution spray at four concentration levels of 2, 4, 6, and 8%, and a control treatment with distilled water, spray, were used. Thus, different liquid solutions were prepared and sprayed onto the plant organs after the plant had grown. Three replications (pots) were considered for each treatment, and other traits were measured except leaf length and width in all three pots.

It is important to note that four repetitions were considered for both leaf length and width. Firstly, the normality test was conducted by Kolmogorov–Smirnov test. In the case of non-normal distribution, a log (10) transformation was used to make them normal. Eight of the 10 traits were normal, and two traits, including flower number and small canopy length, were abnormal.

However, a one-way analysis of variance (ANOVA) was used to analyze the normal and transformed data of traits, and Tukey's test was applied for mean comparisons. Additionally, principal component analysis (PCA) was conducted to identify traits with high variability and sensitivity to different nutritional treatments. The Principal component analysis method has been used in various sciences to determine important parameters in differentiating various phenomena from each other, the results of which include eigenvalues, % of the variance of each variable, factor loading, and broken stick eigenvalue (BSE) ^[26]. All statistical analyses were performed using SPSS software.

Findings

Results of Analysis of Variance

The analysis of variance showed that the effect of different treatments on the characteristics of leaf length, leaf width, number of flowers and fruits, considerable canopy length, small canopy length, plant height, and shoot weight is significant (Table 1). The results showed that the effect of the treatments on the traits of root length and root weight was insignificant.

The mean comparison revealed that

Table 1) The results of analysis of the variance for various traits for *D. stramonium*.

Traits/d.f	Source of Variation	Error	Total
d.f.	8	27	35
Mean Squares			
Leal Length	15.01**	2.74	
Leaf Width	8.25**	1.36	
Traits/d.f	8	18	26
Mean Squares			
Plant Height	68.26*	22.52	
Root Length	17.40 ^{ns}	10.44	
Number of Leaves	143.71 ^{ns}	64.56	
Number of Flowers	150.58**	9.88	
Length of Large Canopy	150.15**	43.08	
Length of Small Canopy	76.83**	12.69	
Weight of Roots	0.32 ^{ns}	0.13	
Weight of Aerial Parts	31.35*	11.71	
Number of Flowers-Log(10)	0.17**	0.03	
Length of Small Canopy Log(10)	0.03**	0.004	

*: Significant at the 0.05 level; **: Significant at the 0.01 level; ns: Not Significant.

the highest plant height (45.75 cm) was observed with the 2% fulvic acid treatment, while the lowest height (31.25 cm) was found in the distilled water treatment. The most significant root length (0.21 cm) was recorded with the 4% fulvic acid treatment, while the lowest (12.75 cm) was associated with the distilled water treatment. For the number of leaves, the highest count (39.50 leaves) was observed in the 4% fulvic acid treatment, while the lowest (19.0 leaves) occurred with the distilled water treatment (Table 2). Regarding the number of flowers, the highest average ^[28] was observed in the 2% fulvic acid treatment, while the lowest (4.5) was found in the 4% solopotas treatment. The distilled water, fulvic acid 8 %, and solopotas 2 % were intermediate levels. In terms of canopy size, the largest

(44.5 cm) and smallest (28.5 cm) canopy lengths were recorded with the 8% fulvic acid treatment, while the smallest (22.5 cm) and largest (15.5 cm) were found with the distilled water treatment (Table 3). For root and shoot weight, the highest values were observed with the 8% solopotas treatment (root weight: 2.65 gr, shoot weight: 13.5 gr), and the lowest values were associated with the distilled water treatment (root weight: 1.65 gr, shoot weight: 3.5 gr). The highest leaf length (14.15 cm) was found with the 2% solopotas treatment, while the lowest (8.17 cm) occurred with the 8% solopotas treatment. For leaf width, the highest (8.67 cm) and lowest (5.07 cm) values were recorded with the 8% fulvic acid and distilled water treatments, respectively (Table 2).

Table 2) Comparison of the means effect of different treatments on the normal traits of *D. stramonium* plant.

Treatment	Plant Height (cm)	Root Length (cm)	Number of Leaves	Length of Large Canopy (cm)	Weight of Roots (gr)	Weight of Aerial Parts (gr)	Leaf Length (cm)	Leaf Width (cm)
Distilled Water	31.25 ±1.75 ^b	12.75 ±3.25 ^a	19.0 ±6.0 ^a	22.5 ±2.5 ^b	1.65 ±0.1 ^a	3.5 ±0.1 ^b	9.25 ±0.9 ^{bc}	5.07 ±1.4 ^d
Fulvic Acid 2 %	45.75 ±1.75 ^a	16.5 ±.5 ^a	33.0 ±2.0 ^a	41.5 ±3.5 ^a	2.3 ±.1 ^a	8.5 ±3.3 ^{ab}	11.37 ±1 ^{bc}	5.62 ±1.0 ^{bcd}
Fulvic Acid 4 %	39.50 ±.50 ^{ab}	21.0 ±2 ^a	39.50 ±15.5 ^a	29.5 ±.50 ^{ab}	1.75 ±0.7 ^a	7.1 ±.4 ^{ab}	10.8 ±1 ^{abc}	6.3 ±1.2 ^{abcd}
Fulvic Acid 6 %	36.50 ±.50 ^{ab}	19.0 ±1 ^a	22.5 ±2.5 ^a	31.50 ±7.5 ^{ab}	2.15 ±0.1 ^a	11.05 ±3.5 ^{ab}	11.6±1 ^{abc}	8.05 ±1.15 ^{abc}
Fulvic Acid 8 %	39.25 ±2.7 ^{ab}	19.0 ±0 ^a	30.50 ±9.5 ^a	44.5 ±7.5 ^a	2.2 ±0.1 ^a	10.6 ±0.1 ^{ab}	12.9 ±0.8 ^{ab}	8.67 ±0.2 ^a
Solopotas 2 %	45.00 ±7 ^a	18.75 ±2.75 ^a	36.0 ±6.0 ^a	39.5 ±10.5 ^{ab}	2.5 ±0.4 ^a	14.15 ±6.65 ^a	14.0 ±3.2 ^a	8.4 ±1.65 ^{ab}
Solopotas 4 %	35.5 ±2.5 ^{ab}	17.25 ±5.75 ^a	27.00 ±7.0 ^a	29.5 ±0.5 ^{ab}	2.1 ±0.2 ^a	9.4 ±1.9 ^{ab}	8.75 ±1.19 ^c	5.4 ±1.3 ^{cd}
Solopotas 6 %	38.5 ±3.5 ^{ab}	18.75 ±1.25 ^a	38.0 ±3.0 ^a	35.0 ±5.0 ^{ab}	2.4 ±0.1 ^a	11.0 ±0.5 ^{ab}	9.65 ±1.0 ^{bc}	5.65 ±1.06 ^{bcd}
Solopotas 8 %	34.00 ±11 ^{ab}	20.0 ±6 ^a	29.5 ±10.5 ^a	29.0 ±11 ^{ab}	2.65 ±0.6 ^a	13.3 ±5.8 ^a	8.17 ±0.9 ^c	5.5 ±0.4 ^{cd}

Table 3) Comparison of the means effect of different treatments on the abnormal traits of *the D. stramonium* plant.

Treatment	Number of Flowers	Number of Flower Log(10)	Length of Small Canopy (cm)	Length of small canopy log(10)
Distilled Water	10.0±3.0 ^b	0.98 ab	15.5±3.5 ^b	1.82b
Fulvic Acid 2 %	28.0±5.0 ^a	1.44a	23.5±5.5 ^{ab}	1.36ab
Fulvic Acid 4 %	8.0±4.0 ^b	0.86b	20.5±0.5 ^{ab}	1.31ab
Fulvic Acid 6 %	5.50±.50 ^b	0.73b	19.0±1.0 ^{ab}	1.27ab
Fulvic Acid 8 %	8.5±2.5 ^b	0.91ab	28.5±7.5 ^a	1.44a
Solopotas 2 %	10.00±1.0 ^b	0.99ab	28.0±2.0 ^a	1.44a
Solopotas 4 %	4.5±2.5 ^b	0.6 b	16.5±0.5 ^b	1.21b
Solopotas 6 %	6.5±0.5 ^b	0.81b	19.0±1.0 ^{ab}	1.27ab
Solopotas 8 %	7.00±5 ^b	0.74 b	15.0±3.0 ^b	1.17b

Principal Component Analysis (PCA) Results

To identify the most important and sensitive traits under the change caused by different treatments and specify the relevant treatment, principal component analysis was used, and the results are presented below (Table 4 and Figure 2). By observing the contribution of each of the components in justifying the traits (Table 3), it can be stated that the traits of considerable crown length, small crown length, plant height, leaf length, leaf width, shoot weight, root length, number of leaves, Root weight and the number of flowers and fruits have the most changes due to different treatments. The results of the principal components analysis showed that 50.9% of the variation in the traits of the *D. stramonium* species was made by the first axis (plant height, length of large canopy, length of small canopy, leaf length, and leaf width) and second axis was made by number of flowers with 20.68 % of the data variance (Table 4).

Table 4) results of analyzing the main components of traits and different treatments.

		Axis (Eigenvector)	
Parameter		1	2
Eigenvalue		5.09	2.06
% Of Variance		50.9	20.68
Cum.% of Variance		50.9	71.59
Broken-stick Eigenvalue		2.92	1.92
Factor Load	Plant Height	-0.859	-0.321
	Root Length	-0.540	0.661
	Number of Leaves	0.621	0.1723
	Number of Flowers	0.272	-0.713
	Length of Large Canopy	0.910	-0.191
	Length of Small Canopy	-0.900	-0.327
	Weight of Roots	0.572	0.523
	Weight of Aerial Parts	-0.662	0.660
	Leaf Length	0.808	-0.364
	Leaf Width	0.736	0.044

Note: Bold parameters are chosen to weigh more in the separation of treatments.

The principal component analysis method was used to extract the most important and sensitive plant traits of *D. stramonium* under different treatments. In this type of analysis, parameters are determined and selected that provide and justify the most information from the available data [27]. To determine the extracted factors and the number of principal components, a large set of indicators such as eigenvalues, scree plots, and percentage of factor load variance are used [27]. This study selected components and factors with a specific value higher than the broken stick eigenvalue and a factor loading of more than 0.5 [28].

By examining the two-plot diagram related to the distribution of treatments in the space of traits, it is clear that almost all plant traits have higher values when solopotas and fulvic acid are used compared to pure water spray. However, in more detail, solopotas spray at a low concentration of about 2% increases leaf width, large crown diameter, and height values. In addition, fulvic acid spray with a concentration of two percent increases the number of flowers and leaf length and causes a relative decrease in root and stem length. To improve this plant's growth efficiency, spraying small amounts of these foods will improve its performance.

Discussion

The growth and performance of plants can be influenced by genetic factors and environmental factors that play the leading role in the occurrence of genetic traits of each plant so that in the absence or deficiency of one of them, the growth of plants will be disturbed and stopped. Proper use of agricultural methods and factors in the production of medicinal plants is important to increase the amount of effective substance. In this regard, the present research was conducted to investigate the effect of different treatments on the structural




Figure 2) Two-plot space of the changes in *D. stramonium* species traits about different treatments using principal component analysis (50.9% and 20.68 % of the data variance are explained by the first and second axes, respectively).

traits and biomass of the medicinal plant *D. stramonium*. The results showed that the effect of different treatments on the characteristics of leaf length, leaf width, number of flowers and fruits, considerable canopy length, small canopy length, plant height, and aerial weight are significant. The comparison of the means showed that the highest height and number of flowers and fruits was related to 2% fulvic treatment, and the lowest amount was related to distilled water treatment. The results of this research are consistent with the results of Arvin (2019) [29], Makkizadeh et al. (2011) [30], Mollashahi et al. (2013) [31] and Hossain et al. (2010) [32]. They reported that when the plant is affected by different concentrations of complete fertilizers, its height traits increase significantly. Potassium has been reported to increase vegetative growth, size, and height in plants by activating enzymes that are effective in growth and

increase the division of meristem cells and the turgescence of meristem cells [33]. This mechanism likely explains the superior performance observed with fulvic acid and potassium treatments in this study. During their research on fennel, they stated that *Azotobacter* inoculation caused a significant increase in the height, side branch, and seed yield of the plant compared to the control [34]. The highest and lowest root length and number of leaves were related to the treatment of 4% fulvic and distilled water, respectively. In research, Arvin (2019) stated that the simultaneous consumption of nutrients nitrogen, phosphorus, and potassium caused an increase in the number of leaves and branches in savory plants compared to the control sample [29]. Researchers also reported that the foliar application of potassium fertilizer increased mung bean area and number of leaves and side branches [35]. In the study of

Alizad et al. (2019), they reported that the combined application of vermicompost with Azospirillum has increased the number of leaves in garlic plants ^[36].

Regarding the traits of considerable canopy length and small canopy length, respectively, in both traits, the highest amount related to fulvic acid treatment was 8%, and the lowest amount was related to a distilled water treatment. The highest amount of root weight and shoot weight traits were related to 8% solopotas treatment, and the lowest amount was related to distilled water treatment. In this regard, the results of Abd Allah et al. (2015) studies showed that salicylic acid foliar application increased the fresh and dry weight of the aerial part of quinoa ^[37]. Additionally, Mosavi Fazl et al. (2015) ^[38] showed that potassium fertilizers enhance dry weight in fodder sorghum, while Habib et al. (2011) ^[39] noted significant increases in wet and dry weights in potato plants following potassium nitrate application. These findings collectively emphasize the crucial role of potassium in biomass accumulation across diverse plant species, including *D. stramonium*. Also, Rahimi and Salahizade (2015) stated that applying potassium sulfate fertilizer in bean plants increased the fresh weight of the stem ^[40]. These results highlight how targeted nutrient application optimizes canopy structure, which is critical for enhancing light capture and growth. Also, Habib et al. (2011) reported that spraying and foliar application of potassium nitrate significantly increased plant length and dry and wet weight in potatoes ^[39]. The highest amount of leaf length is related to 2% solopotas treatment, and the lowest amount is 8% about solopotas treatment. Regarding leaf width, this trait's highest and lowest levels were related to 8% fulvic treatment and distilled water. Therefore, according to the results, using the fertilizers mentioned above

has improved the structural characteristics and biomass of the *D. stramonium* plant. The results of the present research are consistent with the results of the studies of Mohammadpour Vashvaei et al. (2021) ^[12] and Mardi et al. (2022) ^[17]. The results of the survey by Mohammadpour Vashvaei et al. (2021) showed that the highest plant height, the number of branches per plant, the number of leaves per plant, and the dry weight of the leaves of the *D. stramonium* plant belonged to the combination treatment of vermicompost and nanobiomic fertilizers ^[12]. Mardi et al. (2022) reported that the use of fertilizers such as supernitroplus has increased plant height, functional traits, and the number of leaves compared to control (no fertilizer use) in garlic (*Allium sativum* L.) ^[17]. Different fertilizers can increase the plant's ability to absorb organic and inorganic substances from the soil solution and increase plant growth ^[41].

Conclusion

Based on the findings of this study, the application of the aforementioned fertilizers of solopotas 2% and fulvic acid 2% significantly improved the structural characteristics, such as the canopy cover of the *D. stramonium* plant. Shoot and root biomass will increase with the addition of solopotas 6%. However, since the effects on bioactive compounds were not assessed, an increase in biomass cannot be directly linked to a rise in the concentration of effective substances. Further studies are needed to evaluate the impact of these fertilizers on the production of bioactive compounds. It is recommended that future research focus on investigating how different fertilizers affect the concentration of bioactive compounds in *D. stramonium*, in addition to their impact on plant biomass. This will help clarify whether improvements in plant growth translate into higher yields of medicinally valuable substances.

Acknowledgments

The authors would like to thank Urmia University for providing the facilities for this research.

Authors' Contributions

All authors contributed to the manuscript and read and approved its final version.

Funding/Support

Personal funding has been spent to conduct this research.

Ethical Permission

None declared by authors.

Conflict of Interests

The authors declare no conflict of interest.

References:

- Foroghi A. Review of medicinal plants based on antimicrobial properties. *Vet. Res. Bio. Pro.* 2021; 35(1): 2-17.
- Beyk-Khormizi A., Sarafraz-Ardakani M.R., Hosseini Sarghein S., Moshtaghioun S. M., Mousavi-Kouhi S.M., Taghavizadeh Yazdi, M.E. Effect of Organic Fertilizer on the Growth and Physiological Parameters of a Traditional Medicinal Plant under Salinity Stress Conditions. *Horticulture.* 2023; 9(6): 701.
- Da Cunha Honorato A., de Assis R.M.A., Maciel J.F.A., Nohara G.A., de Carvalho AA., Pinto J.E. B.P., Bertolucci S.K.V. Fertilization with different manure sources and doses provides quantitative-qualitative gains in the production of *Thymus vulgaris* L. *S. Afr. J. Bot.* 2024; 164(1): 345-355.
- Hao D., Luan Y., Wang Y., Xiao P. Unveiling Nitrogen Fertilizer in Medicinal Plant Cultivation. *Agron.* 2024; 14(8):1647.
- Ibrahim M.H., Jaafar H.Z., Karimi E., Ghasemzadeh A. Impact of organic and inorganic fertilizers application on the phytochemical and antioxidant activity of Kacip Fatimah (*Labisia pumila* Benth). *Molecules.* 2013 ; 18(9):10973-88.
- Oseni O., Olarinoye C., Amoo I. Studies on chemical compositions and functional properties of thorn apple (*Datura stramonium* L) Solanaceae. *Africa J. Food Sci.* 2011;5(2):40-44.
- Niakan M., Rezapour Mahjoob S., Ghorbanli M. Effect of exogenous putrescine on growth, photosynthesis and alkaloid compounds of *Datura* (*Datura stramonium* L.) in response to salinity stress under hydroponic conditions. *J. Soil Plant Interact.* 2015; 6(1):111-123.
- Al-Snafi A.E. Medical importance of *Datura fastuosa* (syn: *Datura metel*) and *Datura stramonium*-A review. *IOSR J. Pharm.* 2017; 7(2):43-58.
- Singh L.R., Singh O.M. *Datura stramonium*: An overview of its phytochemistry and pharmacognosy. *Res. J. Pharm. Phyto.* 2013; 5(3):143-148.
- Ashtiania F., Sefidkonb F. Tropane alkaloids of *Atropa belladonna* L. and *Atropa acuminata* Royle ex Miers plants. *J. Med. Pla. Res.* 2011; 5(29):6515-6522.
- Heidarzadeh A., Mohammad Modarres-Sanavy S.A., Mokhtassi-Bidgoli A. Effect of nitrogen on some quantitative and qualitative traits of *Dracocephalum kotschy* Boiss. *Iran. J. Hort. Sci.* 2020; 51(3): 705-717.
- Mohammadpour Vashvaei R., Ghanbari A., Asgharipour M.R., Ramroudi M., Dahmardeh M. Evaluation of physiological, biochemical and agronomic characteristics of thorn apple (*Datura stramonium* L.) under residual effect of conventional and low input cropping systems. *Iran. J. Hort. Sci.* 2021; 51(4):995-1015.
- Izadi Z., Biabani A., Sabouri H., Bahreyninejad B. Effects of nitrogen and density on growth and *Datura stramonium* L. yield. *Iran J. Med. Aro. Pla.* 2021; 37(2):315-328.
- Afsharypuor S., Mostajeran A., Shaneh-Saz M. Effect of urea fertilizer on the weight of different parts of *Datura stramonium* and their alkaloidal contents at different developmental stages. *DARU J. Pharm. Sci.* 1997; 7(1):66-72.
- Abbaspour R., Imam Y., Hosseini S.Z. Investigating the effect of organic and chemical fertilizer application on growth and distribution of *Datura* weed seeds. 13th National Congress on Agronomy and Plant Breeding, Karaj, Iran, 2014; 4-6 September: 1-4.
- Hadinezhad F., Salimi A., Chavoushi M. Evaluation of the effect of nitrogen fertilizer and urea fertilizer on some physiological, phytochemical, and antioxidant capacity of *Nigella sativa* L. *Appl. Bio.* 2021; 34(2):168-186.
- Mardi M., Abbasifar A., ValizadehKaji B. Comparison of the effect of biological and non-biological fertilizers on quantitative, qualitative and phytochemical properties of garlic (*Allium sativum* L.). *J. Veg. Sci.* 2022; 5(2):1-17.
- Ehsani S., Niknahad Gharmakher H., Motamedi J., Akbarlou M., Sheidai Karkaj E. Impact of Wheat Straw Biochar and Lignite on Plant Growth of *Astragalus podolobus* Boiss. & Hohen. *ECOPERSIA* 2021; 9(1):61-67.
- Najafzadeh Nobar Z., Safari Sinigani A.A. Effect of soil contamination and antibiotics application on growth and some physiological traits of beans in greenhouse condition. *ECOPERSIA* 2022; 10(4):267-283.
- Abbasi Khalaki M., Ghorbani A., Esmali Ouri A.,

- Shokouhian A., Ashraf Jafari A. Some Facilitators Effects on Alfalfa and Sainfoin Growth in Restoration of Dry-Farming Lands (Study Area: Balekhlichay Watershed, Ardabil, Iran). *ECOPERSIA* 2021; 9(1):43-51.
21. Gorji Z., Mehrparvar M., Mozaffari H. Organic vs. Inorganic Fertilizers: Effects on Growth Parameters and Phenol Content in Cultivated and Wild Safflower. *ECOPERSIA* 2023; 11(2):163-174.
 22. Seilsepour M., Moharami S. Effect of Nitrogen and Municipal Solid Waste Compost on the Physicochemical Properties of a Calcareous Soil under Wheat Cultivation. *Communications in Soil Science and Plant Analysis*. 2024; 1-12.
 23. Zhao Y., Wei Y., Zhang Y., Wen X., Xi B., Zhao X., Wei Z. Roles of composts in soil based on the assessment of humification degree of fulvic acids. *Eco. Indic.* 2017;72(1):473-480.
 24. Patnaik P. *Handbook of inorganic chemicals*: McGraw-Hill New York; 2003. 1086p.
 25. Piccolo A. The Supramolecular structure of humic substances. A novel understanding of humus chemistry and implications in soil science. *Advan. Agron.* 2002; 75(1): 57-134.
 26. Sasanifar S., Alijanpour A., Shafiei Banj A., Eshaghi Rad J.E., Molaei M., Azadi H. Forest protection policy: Lesson learned from Arasbaran biosphere reserve in Northwest Iran. 2019; *La Use Pol.*, 87, 104057.
 27. Hair Junior J.F., Black W.C., Babin B.J., Anderson R.E., Tatham R.L. *Multivariate data analysis*. New Jersey. 1998; 5(3):207-219.
 28. Sheidai-Karkaj E., Sepehry A., Barani H., Motamedi J., Shahbazi F. Establishing a suitable soil quality index for semi-arid rangeland ecosystems in northwest of Iran. *J. Soil Sci. Plant Nutr.* 2019; 19(1): 648-658.
 29. Arvin P. Study of different levels of nitrogen, phosphorus and potassium on physiological and morphological parameters and essential oils in savory plant (*Satureja hortensis* L.). *J. Plant Res.* 2019;32(1):464-473.
 30. Makkizadeh M., Chaichi M., Nasrollahzadeh S., Khavazi K. Effect of different types of nitrogen fertilizers on quantitative and qualitative characteristics of *Satureja hortensis* L. *Iran J. Med. Aro. Plant. Res.* 2012; 28(2):330-341.
 31. Mollashahi M., Ganjali H., Fanaei H. Effect of different levels of nitrogen and potassium on yield, yield components and oil content of sunflower. *Inter. J. Farm. Allie. Sci.* 2013; 2(1):1237-1240.
 32. Hossain M.D., Musa M.H., Talib J., Jol H. Effects of nitrogen, phosphorus and potassium levels on kenaf (*Hibiscus cannabinus* L.) growth and photosynthesis under nutrient solution. *J. Agr. Sci.* 2010; 2(2): 49-57.
 33. Fageria N.K. *The use of nutrients in crop plants*: CRC press; 2016: 448 p.
 34. Tehlan S., Thakral K., Nandal J. Effect of Azotobacter on plant growth and seed yield of fennel (*Foeniculum vulgare* Mill). *Harya J. Hortic. Sci.* 2004; 33(3/4): 287-288.
 35. Thalooth A., Tawfik M., Mohamed H.M. A comparative study on the effect of foliar application of zinc, potassium, and magnesium on growth, yield and some chemical constituents of mungbean plants grown under water stress conditions. *World J. Agri. Sci.* 2006; 2(1):37-46.
 36. Alizad L., Mostafavi Rad M., Aghaei K. Effect of nitrogen sources type and plant growth promoting bacteria on yield and its attributes of Talesh local garlic in Rasht. *J. Crop. Improv.* 2018; 20(2): 533-545.
 37. Abd Allah M.M., El-Bassiouny H.M., Elewa T.A.E., El-Sebai T.N. Effect of salicylic acid and benzoic acid on growth, yield and some biochemical aspects of quinoa plant grown in sandy soil. *Int. J. Chem. Tech. Res.* 2015; 8(1):216-25.
 38. Mousavi Fazl S.H., Alizadeh A., Ansari H., Rezvani Moghaddam P. Effect of different levels of irrigation water and potassium fertilizer on root and shoot growth of forage sorghum. *Iran. J. Irrig. Draig.* 2014; 8(4):747-756.
 39. Habib H.A., Shafeek M., Zaki M., El-Shal Z. Response of potato plants (*Solanum tuberosum* L.) to foliar application with different sources of potassium. *Int. J. Acad. Res.* 2011; 3(1): 129-132.
 40. Rahimi M.M., Salahizadeh A.A. Effect of different levels of irrigation and potassium on qualitative and quantitative characteristics of the beans in Yasooj, Iran. *Eur. J. Natu. Soc. Sci.* 2015; 4(1):50-56.
 41. Lin S., Sattelmacher B., Kutzmutz E., Mühling K.H., Dittert K. Influence of nitrogen nutrition on tuber quality of potato with special reference to the pathway of nitrate transport into tubers. *J. Plant Nutr.* 2004; 27(2):341-350.