

Levels of Mercury in Tissues of Green Tiger Prawn *Penaeus semisulcatus* from Persian Gulf

Mehdi Hosseini^{1*}, Samira Ghiyasi², Mehdi Najafian³ and Abdollah Raeisi Sarasiab⁴

¹ Department of Marine Biology, Faculty of Biological Science, Shahid Beheshti University, Tehran, Iran.

² Department of Environment, Faculty of Agriculture, Islamic Azad University, Arak Branch, Arak, Iran.

³ Department of Environment, Faculty of Natural resources, Islamic Azad University, North Tehran Branch, Tehran, Iran.

⁴ Department of Marine Biology, Faculty of Marine Science, Khoramshahr University of Marine Sciences and Technology, Khoramshahr, Iran.

Received: 21 January 2013 / Accepted: 17 December 2013 / Published Online: 15 June 2014

ABSTRACT Mercury levels in hepatopancreas, muscle and exoskeleton tissues of male and female green tiger prawn, *P. semisulcatus*, from the Persian Gulf were compared. Significantly higher levels ($p < 0.05$) of mercury in female prawn and hepatopancreas tissue were encountered ($p < 0.05$). Significantly higher concentration ($p < 0.05$) of the mercury was also observed in summer than in winter season. This variation could result in internal biological cycle of the organism or variation in bioavailability of mercury in environment.

Key words: Green tiger prawn, Mercury, *Penaeus semisulcatus*, Persian Gulf

1 INTRODUCTION

Historically, the sea has been a major source of protein to the people of Iran and, notwithstanding the variety of imported foods, fishing is still an important occupational and recreational activity today. The fringing reefs, lagoons and offshore waters provide habitats for a great diversity of edible marine life, including a variety of algae, mollusca, crustaceans, sea cucumbers and many different kinds of fish. Local inhabitants commonly harvest representatives from each of these groups for sale or home consumption (Amesbury *et al.*, 1986). Aquatic environments, such as the Persian Gulf, are especially at high risk for mercury contamination since much of

the atmospheric deposition and all of the industrial water-runoffs culminate in these ecosystems.

Mercury is a heavy metal that can be present in the environment in many different forms. For example, mercury can be transformed into a highly toxic compound called methyl mercury, which can accumulate in living organisms and biomagnify (i.e., increase in concentration) as it moves up the food chain. This is the form of mercury to which humans are most often exposed, primarily through consumption of fish and other seafood. For example, fish-eating predators such as loons and larger fish have been observed to bioaccumulate high levels of methyl mercury. Depending on the level of

* Corresponding author: Department of Marine Biology, Faculty of Biological Science, Shahid Beheshti University, Tehran, Iran. Tel: +98 919 528 9331, E-mail: Smhbio@yahoo.com

exposure, effects on humans, fish and wildlife can include slower growth, reproductive failure, and the development of abnormal behaviors that can affect survival. Much of the variation in trace metal tissue concentrations in aquatic organisms has been attributed to the variety in size as well as age of individuals (Farkas *et al.*, 2003), sex (Canli and Furness, 1995; Pourang and Amini, 2001), feeding habits (Kalay *et al.*, 2001; Yilmaz, 2005) and the season of capture (Kargin, 1996; Kargin *et al.*, 2001). These mercury levels in marine environment should be monitored regularly to check water quality, animal health and in view of the quality of public food supplies.

It is known that certain forms of mercury can readily accumulate within crustacean tissues at much higher levels than those in the water column and in sediment (Beltrame and Marco, 2010). More than a decade ago, decapod crustaceans were introduced as biomonitoring organisms for metals in industrially polluted environments due to some suitable characteristics such as their convenient size, abundance, ease of handling in the laboratory and the ability to accumulate metals (Farkas *et al.*, 2003). In terms of commercial catch, green tiger shrimp *P. semisulcatus* remains the most important species in the Persian Gulf, representing 80% of the landed catch in Iran (Niamaimandi *et al.*, 2008).

The Persian Gulf is a body of water in the Middle East between the Arabian Peninsula and Iran. This inland sea is connected to the Gulf of Oman by the Strait of Hormuz (ROPME, 1999). Persian Gulf is a semi-enclosed formation and heavy discharges of the surrounding industries have been ongoing for many decades. Other sources of Persian Gulf pollution include invasions and bombardments that have been staggering in the recent years and are yet to be fully investigated. Large areas of agricultural lands, local fisheries, oil export facilities and a petrochemical plant operate in the general area.

There have been a large number of studies on heavy metals in marine animals recently, particularly in heavily polluted areas of the Persian Gulf (Hosseini *et al.*, 2012; Abdolhpur Monikh *et al.*, 2011; Safaeih *et al.*, 2011; Elahi *et al.*, 2012). In this study, we investigated the relationship between mercury level in tissues (hepatopancreas, muscle, and exoskeleton) with the length and weight of male and female greentiger prawn, *P. semisulcatus*, from the Persian Gulf in summer and winter.

2 MATERIALS AND METHODS

The study was carried out in the several adjoining estuaries in the Persian Gulf (48°25' to 56°25" E, and 24°30' to 30°30' N), including Musa, Bahrekan, Ahmadi and Zangi estuary (Figure 1).

One hundred eighty individuals (90 males and 90 females) of green tiger prawn, *P. semisulcatus*, were collected during summer and winter seasons, 2011.




Figure 1 Map of the Persian Gulf showing sampling stations and study site

Sampling covered areas of the direct or indirect influence of urban and industrial releases, those located near the mouth of the tributary rivers which discharge the industrial pollutants, and a control locality under no industrial or urban releases. The sampling stations were selected to reflect progression of pollution and human activities in the area.

Samples were transferred to the laboratory for further analysis. Each shrimp was properly cleaned by rinsing with distilled water to remove debris, plankton and other external adherent; then, dissected to collect hepatopancreas, muscle and exoskeleton tissues. Then, samples were drained under folds of filter, weighed, wrapped in aluminum foil and then frozen at 10°C prior to analysis. The tissues were placed in clean watch glasses and were oven dried at 105°C for 1 hour and later cooled in a desiccator. Each sample was homogenized in an acid-cleaned mortar and 2 g were digested in triplicate in a water bath at 60°C for 6 h after adding 2.5 mL each of concentrated HNO₃ and H₂SO₄ (Abdolhpur Monikh *et al.*, 2011).

The analysis of total Hg was done by the cold vapor method (Abdolhpur Monikh *et al.*, 2011), using a Perkin-Elmer Atomic Absorption System AA-2380 with automatic background correction and a Perkin-Elmer Mercury Analysis System 303-0830. Replicate (3 to 5) measurements were made on each sample. All the reagents used were of spectroscopic grade and ultra-high purity (99.9%). In order to check for possible contamination, several blanks were performed with the reagents used throughout the course of the experiment. The data obtained were statistically analyzed for confirmation of the results. Mercury from different tissues was calculated by using regression equation and results were expressed in $\mu\text{g g}^{-1}$ dry weight. Statistical analysis was performed using SPSS program. In order to assess significant differences

between tissues, one-way ANOVA and Duncan multiple comparison test were applied. The significance level was set at $\alpha = 0.05$.

3 RESULTS AND DISCUSSION

Mercury level in the tissues of the prawn (*P. semisulcatus*) from four stations in the Persian Gulf is presented in Table 2.

Mercury levels in the tissues of female prawns (0.45 to 4.35 $\mu\text{g g}^{-1}$) were much higher ($P < 0.05$) than in the male (0.39 to 2.02 $\mu\text{g g}^{-1}$). There have been several studies on accumulation of mercury in crustaceans, yet few studies have considered its variation between the sexes. The difference in mercury accumulation between the sexes has been mainly attributed to differences in diet, habitats or due to depuration in eggs, sexual dimorphism and niche partitioning of the forage base (Beckvar *et al.*, 1996). For instance, the male prawn reportedly feeds more on fish and shrimp, while the females feed more on bivalvia, plant and detritus (Williams 1982). Being more in contact with the sediment, plants, bivalves, and detritus receive more sediment associated metals. Gewurtz *et al.* (2011) believed that higher metals levels in female fish were due to the increased consumption of food, relative to males, to meet the increasing demands of reproduction.

Mean mercury level in the hepatopancreas of both sexes was much higher than the other tissue ($P < 0.05$), followed by muscle and exoskeleton, which corresponds with the result of another study (Houserova *et al.*, 2006).

Patterns of mercury accumulations were quite similar for the same sex (Figure 2). Metals are taken up by crustaceans from food and water, distributed throughout crustacean body by blood and eventually accumulated in target organs. Some tissues such as hepatopancreas are considered as one of the main target organs for mercury accumulation (Yilmaz, 2003). The

very high levels of Hg in the hepatopancreas may be related to the content of metallothionein protein in this tissue, which has been found to play a significant role in the regulation and detoxification of mercury (Sen and Semiz,

2007). This protein contains a high percentage of amino group, nitrogen and sulphur that sequester metals in stable complexes (Houserova *et al.*, 2006).

Table 1 Species and specific characteristics

Scientific name	Sex	n	Lenght (cm)	Weight (g)
<i>Penaeus semisulcatus</i>	Male (♂)	90	16.20 ± 0.22	62.03 ± 2.27
	Female (♀)	90	18.10 ± 0.17	76.70 ± 0.80

Mean ± SE

Table 2 Mean concentration of Hg (µg g⁻¹) in tissues of male and female *P. semisulcatus* from Persian Gulf

Season	Tissue	Sex	Station			
			S1	S2	S3	S4
Summer	H	Male	1.20 ± 0.01	2.02 ± 0.02	0.58 ± 0.05	0.98 ± 0.05
		Female	2.30 ± 0.02	4.35 ± 0.01	0.88 ± 0.05	1.23 ± 0.06
	M	Male	1.10 ± 0.03	1.33 ± 0.04	0.75 ± 0.0	0.85 ± 0.05
		Female	1.20 ± 0.04	1.20 ± 0.40	0.88 ± 0.01	0.96 ± 0.02
	E	Male	0.57 ± 0.05	0.95 ± 0.06	0.64 ± 0.02	0.64 ± 0.03
		Female	0.80 ± 0.05	1.03 ± 0.06	0.75 ± 0.02	0.77 ± 0.06
inter	H	Male	1.05 ± 0.04	2.05 ± 0.04	0.51 ± 0.05	0.75 ± 0.01
		Female	1.05 ± 0.04	4.80 ± 0.02	0.72 ± 0.04	0.93 ± 0.02
	M	Male	0.96 ± 0.02	0.95 ± 0.05	0.60 ± 0.02	0.61 ± 0.03
		Female	0.95 ± 0.02	1.90 ± 0.06	0.54 ± 0.01	0.78 ± 0.04
	E	Male	0.60 ± 0.01	0.99 ± 0.04	0.39 ± 0.03	0.56 ± 0.03
		Female	0.65 ± 0.01	1.78 ± 0.02	0.75 ± 0.04	0.45 ± 0.03

H: Hepatopancreas; M: Muscle; E: Exoskeleton




Figure 2 Mercury levels in the all tissues of green tiger prawn *P. semisulcatus* from Persian Gulf

Muscle has also been found to be one of the main target organs for metals accumulation, possibly due to the tendency of metals to react with the sulfhydryl groups of methionine and cysteine proteins that are at high levels in the muscle (Houserova *et al.*, 2006). According to Mieiro *et al.* (2009), in polluted aquatic habitats, hepatopancreas is the main target organ for metals, while in moderately polluted environment metals are accumulated in muscle. The absence or low value of metals level in some tissues may indicate that those tissues are not the target organs for metals accumulation or may be due to the major functional differences in their body (Sen and Semiz, 2007).

Significantly higher ($P < 0.05$) mercury level was also measured in summer than in winter for both male ($2.02 \mu\text{g g}^{-1}$) and female prawns ($4.35 \mu\text{g g}^{-1}$). This variation could result in internal biological cycle in organism or variation in bioavailability of metal in environment. Temperature, food availability and water could increase metal concentration in summer more than in winter (Sen and Semiz, 2007). Higher mercury level in invertebrates during summer has also been reported earlier (Bordin *et al.*, 1992; Hosseini *et al.*, 2012), which has been attributed to its accumulation in fatty tissues that sharply increase during the reproduction in summer.

Since larger organisms at higher trophic levels generally exhibit higher contaminant level in their bodies (Abdolapur Monikh *et al.*, 2012), prawn that feed on larger food items also accumulate more contaminants as compared with the crustaceans that eat smaller organisms. We expected to see higher mercury levels in tissues of female prawn because they are larger and can eat larger food items.

There was a significant difference ($P < 0.05$) in mercury level among different stations, the highest being recorded in the samples from Musa estuary and the least was from Ahmadi estuary during both seasons. The high mercury levels in the Musa estuary can be explained by

its vicinity to Imam port with its tremendous oil tankers traffics, enormous discharges of petrochemical units, agricultural and urban development activities in the area (Safahieh *et al.*, 2011; Abdolapur Monikh *et al.*, 2012). Crustaceans collected from the Bahrekan estuary also demonstrated relatively high metals levels in their tissues in comparison to those from Ahmadi and Zangi estuaries. Bahrekan estuary is used for many industrial, shipping and irrigation purposes. In addition, it passes several urban and rural areas, ports and harbors.

4 CONCLUSION

Current study provides new information on the distribution of mercury in the tissues of green tiger prawn *P. semisulcatus* from the Persian Gulf. In general, the mercury showed a strong tendency to accumulate in the hepatopancreas, followed by muscle and exoskeleton tissues, respectively. A higher level of mercury in the tissues of female prawn was also recorded. The highest concentrations of mercury in the tissues were recorded in Musa estuary. Mercury levels in the edible part of the prawn are below the proposed limit values for human consumption, except in some cases.

5 ACKNOWLEDGEMENT

The author would like to thank Prof. Jabar Salamat, Moemen Ali Peery, Akbar Almasi and Dr. Ali Hossein Jalilian. This research was supported by the Environment Protection Institute.

6 REFERENCES

- Abdolapur Monikh, F., Peery, S., Karami, O., Hosseini, M., Bastami, A.A. and Ghasemi, A.F. Distribution of Metals in the Tissues of Benthic, *Euryglossa orientalis* and *Cynoglossus aereus*, and Benthic-Pelagic, *Johnius belangerii*, Fish from Three

- Estuaries, Persian Gulf. Bull. Environ. ContamToxicol., 2012; 18:319-324.
- Bordin, G.Mc-Court, J. and Rodriquez, A. Trace metal in the marine bivalve *Macomabalthica* in the westerschelde estuary the Nerthlands. Part 1: Analysis of total copper, cadmium, zinc and iron concentrations; local and seasonal variations. Science of the Total Environ., 1992; 127 (3): 255-280.
- Birke, M. and Rauch, U. Urban geochemistry: Investigation in the Berlin metropolitan area. Environ. Geochem. Hlth., 2000; 22: 233-248.
- Boyden, C.R. and Phillips, D.J H. Seasonal variation and inherent variability of trace elements in oysters and their implications for indicator studies. Mar. Ecol-Prog. Ser., 1981; 5: 29-40.
- Canli, M. and Furness, R.W. Toxicity of heavy metals dissolved in seawater and influences of sex and size on metal accumulation and tissue distribution in the Norway lobster *Nephropsnorvegicus*. Mar. Environ. Res., 1993; 36: 217-236.
- El-Moselhy, K.H.M. Bioaccumulation of mercury in some marine organisms from Lake Timsah and Bitter Lakes, Suez Canal, Egypt. Egyption J. Aquatic Res., 2006; 1: 124-134.
- FAO, Committee for inland fisheries of Africa: Report of the third session of the working party on pollution and fisheries, FAO Fish. Rep., 1992; 471: 43P.
- Falusi, B.A. and Olanipekun, E.O. Bioconcentration factors of heavy metals in tropical crab (*Carcinussp*) from River Aponwe, Ado-Ekiti, Nigeria. J. Appl. Sci. Environ. Manage., 2007;11(4): 51-54.
- Farkas, A. Salanki, J. and Specziar, A. Age and size-specific patterns of heavy metals in the organs of freshwater fish *Abramisbrama* L. populating a low-contaminated site. Water Res., 2003; 37: 959-964.
- Hosseini, M. Nabavi, S.M.B. Bastami, A.A. and Parsa, Y. Mercury Concentration in Tissues of Blue Swimming Crab, *Portunuspelagicus* (Linnaeus, 1758) and Sediments from Persian Gulf Coasts, Iran. World Appl. Sci. J., 2012; 18 (3): 322-327.
- Kargin, F. Donmez, A. and Cogun, H.Y. Distribution of heavy metals in different tissues of the shrimp *Penaeus semiculatus* and *Metapenaeus monocerus* from the Iskenderun Gulf, Turkey: seasonal variations, B. Environ. Contam. Tox., 2001; 66: 102-109.
- Kargin, F. Seasonal changes in levels of heavy metals in tissues of *Mullusbarbatus* and *Sparusaurata* collected from Iskenderun Gulf (Turkey). Water Air Soil Poll., 1996 90: 557-562.
- Kumar, M. Ferguson, G. Xiao, Y. Hooper, G. and Venema, S. Studies on reproductive biology and distribution of blue swimmer crab (*Portunuspelagicus*) in South Australian Waters, Res. report series., 2000; 47: 1324-2083.
- Landner, L. and Lindestrom, L. Zinc in society and in the environment, Miljoforskargruppen, Stockholm. 1998.
- Niamaimandi, N., Azizi, A., Khalijah, D.S., Roos, S.C. and Kiabi, B. Reproductive biology of the green tiger prawn (*Penaeus semisulcatus*) in coastal waters of Bushehr, Persian gulf. ICES J. Mar. Sci., 2008; 65:1593-1599.

- Pourang, N. and Amini, G. Distribution of trace elements in tissues of two shrimp species from Persian Gulf and effects of storage temperature on elements transportation. *Water, Air, Soil Poll.*, 2001; 129: 229-243.
- Safahieh, A. Abdolahpur Monikh, F. and Savari, A. Heavy metals contamination in sediment and Sole fish (*Euryglossa orientalis*) from Musa estuary (Persian Gulf). *W J. Fish Mar. Sci.*, 2011; 3:290-297.
- Storelli, M.M. Storelli, A.D. Ddabbo, R. Marano, C. Bruno, R. and Marcotrigiano, G.O. Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea. Overview and evaluation. *Environ. Pollut.*, 2005; 135: 163-170.
- Reinecke, A.J. Snyman, R.G. and Nel, J.A.J. Uptake and distribution of lead (Pb) and cadmium (Cd) in the freshwater crab, *Potamonautes perlatus* (Crustacea) in the Eerste River, South Africa. *Water Air Soil Poll.*, 2003; 145: 395- 408.
- Valinassab, T. Daryanabard, R. Dehghani, R. and Pierce, G.J. Abundance of demersal fish resources in the Persian Gulf and Oman Sea. *J. Mar. Biol. Assoc. UK.*, 2006; 86:1455-1462.
- Yi, Y. Wang, Z. Zhang, K. Yu, G. and Duan, X. Sediment pollution and its effect on fish through food chain in the Yangtze River. *Int. J. Sediment Res.*, 2008; 23:338-347.
- Yilmaz, F., Ozdemir, N., Demirak, A. and Levent, T.A. Heavy metal levels in two fish species *Leuciscus cephalus* and *Lepomis gibbosus*. *Food Chem.*, 2007; 100: 830-835.
- Yilmaz, A.B. and Yilmaz, L. Influences of sex and seasons on levels of heavy metals in tissues of green tiger shrimp (*Penaeus semisulcatus*). *Food Chem.*, 2007; 101: 1664-1669.

غلظت جیوه در بافت‌های میگوی ببری سبز *Penaues semisulcatus* در سواحل خلیج فارس

مهدی حسینی^۱ سمیرا قیاسی^۲ مهدی نجفیان^۳ عبدالله رئیسی سراسیاب^۴

- ۱- گروه زیست‌شناسی دریا، دانشکده علوم زیستی، دانشگاه شهید بهشتی، تهران، ایران
- ۲- گروه محیط زیست، دانشکده کشاورزی، دانشگاه آزاد اسلامی واحد اراک، اراک، ایران
- ۳- گروه محیط زیست، دانشکده منابع طبیعی، دانشگاه آزاد اسلامی واحد تهران شمال، تهران، ایران
- ۴- گروه زیست‌شناسی دریا، دانشکده علوم دریایی، دانشگاه علوم و فنون دریایی خرمشهر، خرمشهر، ایران

تاریخ دریافت: ۱ بهمن ۱۳۹۱ / تاریخ پذیرش: ۲۶ آذر ۱۳۹۲ / تاریخ چاپ: ۲۵ خرداد ۱۳۹۳

چکیده غلظت جیوه در بافت‌های هیپاتوپانکراس، ماهیچه و اسکلت خارجی میگوی ببری سبز *P. semisulcatus* در سواحل خلیج فارس بررسی شد. تغییرات غلظت جیوه با جنس و فصل ارتباط داشت. نتایج نشان داد که غلظت جیوه در هیپاتوپانکراس نسبت به ماهیچه و اسکلت خارجی بیشتر است. اختلاف معنی‌داری بین غلظت جیوه در میگوی نر و ماده وجود داشت، به طوریکه جنس ماده غلظت بالاتری داشت. همچنین میگوی ماده نزدیک بستر زیست می‌ند و جیوه‌ای که با بستر تماس دارد را دریافت می‌کند. نتایج همچنین نشان داد که جیوه در بافت‌های میگو در فصل تابستان نسبت به فصل زمستان غلظت بالاتری داشت. تغییرات فصلی ممکن است با چرخه زیستی در میگو و میزان در دسترس بودن فلز در محیط ارتباط داشته باشد.

کلمات کلیدی: *Penaues semisulcatus*، جیوه، خلیج فارس، میگوی ببری سبز